



Cochrane
Library

Cochrane Database of Systematic Reviews

Interprofessional collaboration to improve professional practice and healthcare outcomes (Review)

Reeves S, Pelone F, Harrison R, Goldman J, Zwarenstein M

Reeves S, Pelone F, Harrison R, Goldman J, Zwarenstein M.

Interprofessional collaboration to improve professional practice and healthcare outcomes.

Cochrane Database of Systematic Reviews 2017, Issue 6. Art. No.: CD000072.

DOI: 10.1002/14651858.CD000072.pub3.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	4
BACKGROUND	6
OBJECTIVES	7
METHODS	7
RESULTS	9
Figure 1.	10
Figure 2.	13
Figure 3.	14
ADDITIONAL SUMMARY OF FINDINGS	18
DISCUSSION	19
AUTHORS' CONCLUSIONS	21
ACKNOWLEDGEMENTS	22
REFERENCES	22
CHARACTERISTICS OF STUDIES	25
DATA AND ANALYSES	44
APPENDICES	44
WHAT'S NEW	45
HISTORY	46
CONTRIBUTIONS OF AUTHORS	46
DECLARATIONS OF INTEREST	47
SOURCES OF SUPPORT	47
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	47
INDEX TERMS	47

[Intervention Review]

Interprofessional collaboration to improve professional practice and healthcare outcomes

Scott Reeves¹, Ferruccio Pelone¹, Reema Harrison², Joanne Goldman³, Merrick Zwarenstein⁴

¹Faculty of Health, Social Care and Education, Kingston University and St George's, University of London, London, UK. ²University of Sydney, Sydney, Australia. ³Centre for Quality Improvement and Patient Safety, University of Toronto, Toronto, Canada. ⁴Department of Family Medicine, University of Western Ontario, London, Canada

Contact address: Scott Reeves, Faculty of Health, Social Care and Education, Kingston University and St George's, University of London, St George's Hospital, Grosvenor Wing, Cranmer Terrace, London, Greater London, SW17 0BE, UK. s.reeves@sgul.kingston.ac.uk.

Editorial group: Cochrane Effective Practice and Organisation of Care Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 6, 2017.

Citation: Reeves S, Pelone F, Harrison R, Goldman J, Zwarenstein M. Interprofessional collaboration to improve professional practice and healthcare outcomes. *Cochrane Database of Systematic Reviews* 2017, Issue 6. Art. No.: CD000072. DOI: 10.1002/14651858.CD000072.pub3.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Poor interprofessional collaboration (IPC) can adversely affect the delivery of health services and patient care. Interventions that address IPC problems have the potential to improve professional practice and healthcare outcomes.

Objectives

To assess the impact of practice-based interventions designed to improve interprofessional collaboration (IPC) amongst health and social care professionals, compared to usual care or to an alternative intervention, on at least one of the following primary outcomes: patient health outcomes, clinical process or efficiency outcomes or secondary outcomes (collaborative behaviour).

Search methods

We searched CENTRAL (2015, issue 11), MEDLINE, CINAHL, ClinicalTrials.gov and WHO International Clinical Trials Registry Platform to November 2015. We handsearched relevant interprofessional journals to November 2015, and reviewed the reference lists of the included studies.

Selection criteria

We included randomised trials of practice-based IPC interventions involving health and social care professionals compared to usual care or to an alternative intervention.

Data collection and analysis

Two review authors independently assessed the eligibility of each potentially relevant study. We extracted data from the included studies and assessed the risk of bias of each study. We were unable to perform a meta-analysis of study outcomes, given the small number of included studies and their heterogeneity in clinical settings, interventions and outcomes. Consequently, we summarised the study data and presented the results in a narrative format to report study methods, outcomes, impact and certainty of the evidence.

Main results

We included nine studies in total (6540 participants); six cluster-randomised trials and three individual randomised trials (1 study randomised clinicians, 1 randomised patients, and 1 randomised clinicians and patients). All studies were conducted in high-income countries (Australia, Belgium, Sweden, UK and USA) across primary, secondary, tertiary and community care settings and had a follow-up of up to 12 months. Eight studies compared an IPC intervention with usual care and evaluated the effects of different practice-based IPC interventions: externally facilitated interprofessional activities (e.g. team action planning; 4 studies), interprofessional rounds (2 studies), interprofessional meetings (1 study), and interprofessional checklists (1 study). One study compared one type of interprofessional meeting with another type of interprofessional meeting. We assessed four studies to be at high risk of attrition bias and an equal number of studies to be at high risk of detection bias.

For studies comparing an IPC intervention with usual care, functional status in stroke patients may be slightly improved by externally facilitated interprofessional activities (1 study, 464 participants, low-certainty evidence). We are uncertain whether patient-assessed quality of care (1 study, 1185 participants), continuity of care (1 study, 464 participants) or collaborative working (4 studies, 1936 participants) are improved by externally facilitated interprofessional activities, as we graded the evidence as very low-certainty for these outcomes. Healthcare professionals' adherence to recommended practices may be slightly improved with externally facilitated interprofessional activities or interprofessional meetings (3 studies, 2576 participants, low certainty evidence). The use of healthcare resources may be slightly improved by externally facilitated interprofessional activities, interprofessional checklists and rounds (4 studies, 1679 participants, low-certainty evidence). None of the included studies reported on patient mortality, morbidity or complication rates.

Compared to multidisciplinary audio conferencing, multidisciplinary video conferencing may reduce the average length of treatment and may reduce the number of multidisciplinary conferences needed per patient and the patient length of stay. There was little or no difference between these interventions in the number of communications between health professionals (1 study, 100 participants; low-certainty evidence).

Authors' conclusions

Given that the certainty of evidence from the included studies was judged to be low to very low, there is not sufficient evidence to draw clear conclusions on the effects of IPC interventions. Nevertheless, due to the difficulties health professionals encounter when collaborating in clinical practice, it is encouraging that research on the number of interventions to improve IPC has increased since this review was last updated. While this field is developing, further rigorous, mixed-method studies are required. Future studies should focus on longer acclimatisation periods before evaluating newly implemented IPC interventions, and use longer follow-up to generate a more informed understanding of the effects of IPC on clinical practice.

PLAIN LANGUAGE SUMMARY

How effective are strategies to improve the way health and social care professional groups work together?

What is the aim of this review?

The aim of this Cochrane Review was to find out whether strategies to improve interprofessional collaboration (the process by which different health and social care professional groups work together), can positively impact the delivery of care to patients. Cochrane researchers collected and analysed all relevant studies to answer this question, and found nine studies with 5540 participants.

Key messages

Strategies to improve interprofessional collaboration between health and social care professionals may slightly improve patient functional status, professionals' adherence to recommended practices, and the use of healthcare resources. Due to the lack of clear evidence, we are uncertain whether the strategies improved patient-assessed quality of care, continuity of care, or collaborative working.

What was studied in this review?

The extent to which different health and social care professionals work well together affects the quality of the care that they provide. If there are problems in how these professionals communicate and interact with each other, this can lead to problems in patient care. Interprofessional collaboration practice-based interventions are strategies that are put into place in healthcare settings to improve interactions and work processes between two or more types of healthcare professionals. This review studied different interprofessional collaboration interventions, compared to usual care or an alternative intervention, to see if they improved patient care or collaboration.

What are the main results of the review?

The review authors found nine relevant studies across primary, secondary, tertiary and community care settings. All studies were conducted in high-income countries (Australia, Belgium, Sweden, UK and USA) and lasted for up to 12 months. Most of the studies were well conducted, although some studies reported that many participants dropped out. The studies evaluated different methods of interprofessional collaboration, namely externally facilitated interprofessional activities (e.g. collaborative planning/reflection activities led by an individual who is not part of the group/team), interprofessional rounds, interprofessional meetings, and interprofessional checklists.

Externally facilitated interprofessional activities may slightly improve patient functional status and health care professionals' adherence to recommended practices, and may slightly improve use of healthcare resources. We are uncertain whether externally facilitated interprofessional activities improve patient-assessed quality of care, continuity of care, or collaborative working behaviours. The use of interprofessional rounds and interprofessional checklists may slightly improve the use of healthcare resources. Interprofessional meetings may slightly improve adherence to recommended practices, and may slightly improve use of healthcare resources.

Further research is needed, including studies testing the interventions at scale to develop a better understanding of the range of possible interventions and their effectiveness, how they affect interprofessional collaboration and lead to changes in care and patient health outcomes, and in what circumstances such interventions may be most useful.

How up to date is this review?

The review authors searched for studies that had been published to November 2015.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Effects of practice-based interprofessional collaboration (IPC) interventions on professional practice and healthcare outcomes compared to usual care

Patient or population: health and social care professionals involved in the delivery of health services and patient care

Settings: primary, secondary, tertiary and community care settings, primarily in the USA and the UK

Intervention: practice-based interprofessional collaboration (IPC) interventions with an explicit objective of improving collaboration between two or more health or social care professionals

Comparison: usual care

Outcomes	Impacts	No. of studies (participants)	Certainty of the evidence (GRADE)
Patient health outcomes			
Patient functional status	Externally facilitated inter-professional activities may slightly improve stroke patients' functional status (Strasser 2008).	1 (464)	⊕⊕⊕⊕ Low^a
Patient-assessed quality of care	It is uncertain if externally facilitated interprofessional activities increases patient-assessed quality of care because the certainty of this evidence is very low (Black 2013).	1 (1185)	⊕⊖⊖⊖ Very low^b
Patient mortality, morbidity or complication rates	None of the included studies reported patient mortality, morbidity or complication rates	--	--
Clinical process or efficiency outcomes			
Adherence to recommended practices	The use of interprofessional activities with an external facilitator or interprofessional meetings may slightly improve adherence to recommended practices and prescription of drugs (Cheater 2005 ; Deneckere 2013 ; Schmidt 1998).	3 (2576)	⊕⊕⊖⊖ Low^c

Continuity of care	It is uncertain if externally facilitated interprofessional activities improves continuity of care because the certainty of this evidence is very low (Strasser 2008).	1 (464)	⊕⊕⊕⊕ Very low^d
^d Use of healthcare resources	Interprofessional checklists (Calland 2011), interprofessional rounds (Curley 1998; Wild 2004) or externally facilitated interprofessional activities (Strasser 2008), may slightly improve overall use of resources, length of hospital stay, or costs	4 (1679)	⊕⊕⊕⊕ Low^e

Collaborative behaviour outcomes

Collaborative working; team communication; team co-ordination	It is uncertain whether externally facilitated interprofessional activities (Black 2013; Calland 2011; Cheater 2005; Deneckere 2013) improve collaborative working, team communication, and co-ordination because the certainty of this evidence is very low	4 (1936)	⊕⊕⊕⊕ Very low^f
---	--	----------	-------------------------------------

GRADE Working Group grades of evidence

High-certainty: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate-certainty: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low-certainty: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very-certainty: We are very uncertain about the estimate.

^a We assessed the certainty of the evidence as low because of high risk of bias (no blinding of outcome assessment).

^b We assessed the certainty of the evidence as very low because of the risk of bias (high risk of attrition and detection bias; details about allocation sequence generation and concealment were not reported).

^c We assessed the certainty of the evidence as low due to potential indirectness (both studies were conducted in one country and the outcomes may not be transferable to other settings), and risk of bias (high risk of attrition, unclear selection and reporting risk).

^d We assessed the certainty of the evidence as very low because of risk of bias (high risk of attrition and detection bias, and unclear risk of selection bias).

^e We assessed the certainty of evidence as low because of high risk of bias (attrition and detection), and unclear risk of bias (selection, reporting, and contamination).

^f We assessed the certainty of the evidence as very low due to high risk of bias (selection, attrition, and detection) or unclear risk of bias (reporting and contamination).

BACKGROUND

Description of the condition

Interprofessional collaboration (IPC) is the process by which different health and social care professional groups work together to positively impact care. IPC involves regular negotiation and interaction between professionals, which values the expertise and contributions that various healthcare professionals bring to patient care. However, IPC can be affected by problems linked to imbalances of authority, limited understanding of others' roles and responsibilities, and professional boundary friction when delivering patient care (Baker 2011; Reeves 2010).

Research has repeatedly documented the impact of collaboration problems on work processes and patient safety (Lillebo 2015; Van Leijen-Zeelenberg 2015). For example, failures of collaboration were found to be at the centre of a number of care failures across the globe (Francis 2013; The Joint Commission 2016). Therefore, professionals must ensure that they collaborate in an effective manner to deliver safe, high-quality patient care.

Different health policy makers across the globe have repeatedly called for the use of IPC as a key approach to improve the quality and safety of patient care (Department of Health 1997; Health Canada 2003; Institute of Medicine 2000; Institute of Medicine 2013; WHO 1976; WHO 2010). During the past 10 years in particular, IPC has been at the forefront of much curricular, research, policy, and regulatory activity at national and international levels. The promotion of IPC stems from the complexity and multifaceted nature of patients' health and care needs and the health system, and research that suggests that improved collaboration between multiple professionals may be essential for the provision of effective and comprehensive care.

Research in the area of IPC is complicated by the use of varied terms (interdisciplinary collaboration, multidisciplinary co-ordination, transprofessional teamwork), which has resulted in conceptual confusion within the field (Reeves 2010). As a result, one must take care when evaluating such studies, to ensure one understands the nature and key activities of the intervention, whatever it may be named.

Description of the intervention

An IPC intervention involves members of more than one health or social care profession interacting together with the explicit purpose of improving IPC. A scoping review examining the nature

of interventions used in the interprofessional field found three main types: education-based interventions, practice-based interventions, and organisationally-based interventions (Reeves 2011). This review focuses on interprofessional practice-based interventions, also called practice-based IPC interventions. An interprofessional practice-based intervention involves the deployment in the workplace of a tool or routine to improve IPC; examples include communication tools, interprofessional meetings, and checklists. A review focusing solely on interprofessional education (an education-based intervention) was recently updated (Reeves 2013). In this review, an interprofessional education intervention was defined as 'members of more than one health or social care profession learning interactively together, for the explicit purpose of improving interprofessional collaboration, the health and well-being of patients, or both.' Interactive learning requires active learner participation and active exchange between learners from different professions.

An interprofessional organisationally-based intervention involves a change at an organisational level to improve interprofessional collaboration; examples include policy and staffing changes (Reeves 2010). A review of the effects of this type of intervention still needs to be undertaken, to generate a more holistic understanding of the nature of these different, but complementary, interventions.

How the intervention might work

A practice-based IPC intervention might work by incorporating a tool, routine, or activity to improve interprofessional interaction (e.g. communication, co-ordination) into clinical practice. In turn, this may improve how healthcare professionals work together and deliver health care, leading to improved health outcomes.

Why it is important to do this review

Research identifying various problems with IPC, and the delivery of care and patient outcomes, continues to accumulate (Körner 2016; Van Leijen-Zeelenberg 2015). Therefore, it is important to understand the effectiveness of interventions aimed at improving IPC on health and social care. Governments around the globe continue to institute major changes and invest significant resources to improve IPC. Ideally, these policy decisions should be based on evidence of the effectiveness of these approaches. The aim of this review is to update a previous review, and synthesise evidence from randomised trials of practice-based IPC interventions, to inform such decision-making (Zwarenstein 2009).

OBJECTIVES

To assess the impact of practice-based interventions designed to improve IPC amongst health and social care professionals, compared to usual care or to an alternative intervention, on at least one of the following primary outcomes: patient health outcomes, clinical process or efficiency outcomes or secondary outcomes (collaborative behaviour).

METHODS

Criteria for considering studies for this review

Types of studies

We only considered individual or cluster-randomised, which provide the most reliable evidence for the effects of practice-based interprofessional collaboration (IPC).

Types of participants

We included interventions that targeted any type of health and social care professional (e.g. chiropodists or podiatrists, complementary therapists, dentists, dietitians, doctors or physicians, hygienists, midwives, nurses, occupational therapists, pharmacists, physiotherapists, psychologists, psychotherapists, radiographers, social workers, or speech therapists).

Types of interventions

We included any practice-based intervention with an explicit objective of improving collaboration between two or more health or social care professionals. We used the following criterion to include interventions.

- Evaluations of a practice-based IPC intervention, where the study explicitly noted an objective to improve collaboration amongst two or more types of health or social care professionals. Other terms besides IPC could have been used, and were accepted as equivalent to IPC, such as communication, co-ordination, and teamwork.

The comparator was usual care or an alternative intervention. We placed no restrictions on interventions or settings (e.g. hospitals, primary care, community-based care).

Types of outcome measures

Primary outcomes

- Patient health outcomes (objectively measured or self-reported, using a validated instrument)
 - mortality
 - morbidity
 - disease incidence
 - disease duration
 - cure rates
 - quality of life measures
 - functional status
 - complication rate
 - patient-assessed quality of care
- Clinical process or efficiency outcomes
 - readmission rates
 - adherence to recommended practices (by healthcare providers)
 - continuity of care
 - use of healthcare resources (i.e. cost-benefit analysis)
 - participant satisfaction

Secondary outcomes

- Collaborative behaviour (objective or self-reported outcomes, using a validated instrument)

We excluded interprofessional learning (interprofessional education) as a secondary outcome. Whilst interprofessional education can support IPC in the workplace, these are distinct activities and our focus was the latter.

Search methods for identification of studies

Electronic searches

We searched the following sources.

1. Cochrane Central Register of Controlled Trials (CENTRAL; 2015, issue 11) in the Cochrane Library (searched on 24 November 2015; full strategy available in [Appendix 1](#)).
2. MEDLINE Ovid: 2007 to 2015 (searched on 10 November 2015; full strategy available in [Appendix 2](#)).
3. CINAHL EBSCO: 2007 to 2015 (searched on 10 November 2015; full strategy available in [Appendix 3](#)).
4. [ClinicalTrials.gov](#) and WHO International Clinical Trials Registry Platform (ICTRP): 2007 to 2015 (searched on 24 November 2015; full strategy available in [Appendix 4](#)).

We placed no language restrictions on the search strategy. We did not search Embase, as a review of the studies included previously showed that none were indexed in this database. We included all the trials identified by the previous version of the review ([Zwarenstein 2009](#)).

Searching other resources

We handsearched the *Journal of Interprofessional Care* (2007 to November 2015), and reviewed the reference lists of included studies.

Data collection and analysis

Selection of studies

At least two review authors (SR, FP) independently reviewed each of the titles and abstracts retrieved in the searches, to identify those that met the review's inclusion criteria.

We obtained the full-text of all potentially relevant articles. At least two review authors (SR, FP) independently assessed each full-text article to determine if it met all of the criteria. We resolved disagreements by consultation with another review author (MZ). As a further quality check, this additional review author reviewed all included articles.

Data extraction and management

Two review authors (SR, FP) independently extracted the following information from included studies.

1. Study setting (country, healthcare setting).
2. Types of study participants.
3. Description of IPC intervention.
4. Description of any other interventions.
5. Outcomes (primary and secondary).
6. Data for the main outcomes.

Assessment of risk of bias in included studies

We used the suggested criteria recommended by Cochrane Effective Practice and Organisation of Care (EPOC) to assess risk of bias in all studies included in the review (EPOC 2016), an approach that assessed the key areas of:

1. selection bias;
2. performance bias;
3. detection bias;
4. attrition bias;
5. reporting bias; and
6. any other potential sources of bias.

For each criterion, we described the relevant information provided by the trial authors, and judged each item as being at: 1) high risk of bias (plausible bias that seriously weakens confidence in the results); 2) low risk of bias (plausible bias unlikely to seriously alter the results); or 3) unclear risk of bias (lack of information or uncertainty over the potential for bias).

We reported all included studies in the [Risk of bias in included studies](#) section below. We did not exclude studies on the basis of their risk of bias. We made an overall assessment of the risk of bias

for each outcome (across criteria) within and across studies, using the approach suggested by EPOC 2016.

Measures of treatment effect

We had initially planned to conduct a meta-analysis, however, this was not possible due to differences in populations and interventions (see [Results](#)). Therefore, we presented a narrative summary of the results.

Unit of analysis issues

We critically examined the methods of analysis of all study types. We identified cluster-randomised trials, and where appropriate, commented on unit of analysis errors in the results and discussion.

Dealing with missing data

It was not possible to undertake a meta-analysis of the included studies due to heterogeneity and therefore the issue of missing data in statistical analysis did not arise.

Assessment of heterogeneity

We did not assess heterogeneity statistically. Our narrative of randomised trials compared the characteristics of the study populations, interventions, and outcomes (see [Data synthesis](#)).

Assessment of reporting biases

We attempted to reduce the risk of reporting bias by undertaking comprehensive searches of multiple databases and trial registers. We found too few studies reporting the primary outcomes to allow any assessment of reporting bias.

Data synthesis

We could not complete a meta-analysis of study outcomes for this review due to the small number of included studies, and the differences in IPC subtypes, settings, participants, and outcomes across the studies. Consequently, we presented the results in a narrative format. In producing this narrative, we grouped (or categorised) the following types of practice-based IPC interventions as: externally facilitated interprofessional activities, interprofessional rounds, interprofessional meetings, and interprofessional checklists.

Two review authors (SR, MZ) independently assessed the certainty of the evidence (high, moderate, low, and very low) as it related to the main outcomes, using the five GRADE considerations (risk of bias, consistency of effect, imprecision, indirectness, and publication bias) Guyatt 2008.

We developed two 'Summary of findings' tables for the comparisons: (1) practice-based IPC compared with usual care and; (2)

practice-based IPC compared with an alternative practice-based IPC. We included the following outcomes.

- Patient health outcomes: patient functional status; patient-assessed quality of care; patient mortality, morbidity or complication rates.
- Clinical process or efficiency outcomes: adherence to recommended practices; continuity of care; use of healthcare resources.
- Collaborative behaviour outcomes: collaborative working; team communication or co-ordination.

We applied the methods and recommendations described in Section 8.5 and Chapter 12 of the *Cochrane Handbook of Systematic Reviews of Interventions* (Higgins 2011), and the EPOC worksheets (EPOC 2013). We justified all decisions to downgrade or upgrade the certainty of the evidence using footnotes, and made comments to aid readers' understanding of the review where necessary.

Subgroup analysis and investigation of heterogeneity

As we did not undertake a meta-analysis, we could not conduct a subgroup analysis.

Sensitivity analysis

We had planned to perform a sensitivity analysis for pooled results based on the risk of bias information recorded from the included studies. However, a meta-analysis was not possible, due to a variation in the intervention and study methods used in the included studies.

RESULTS


Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

Results of the search

Our searches generated a total of 2493 abstracts. Once duplicate abstracts were removed (N = 1083), the total number of abstracts reviewed was 1410 (see [Figure 1](#)). The handsearch did not produce any additional articles. Following assessment of each of the abstracts, we identified 34 studies that potentially met our inclusion criteria (1 from EPOC, 1 from CENTRAL, 29 from MEDLINE, and 3 from CINAHL).

Figure 1. Flow diagram of study selection



We included nine studies (N = 6540), all conducted in high-income countries (Australia, Belgium, Sweden, UK, and USA). We identified four studies for this update ([Black 2013](#); [Calland 2011](#); [Deneckere 2013](#); [Strasser 2008](#)).

Eight studies compared an interprofessional collaboration (IPC) intervention to a control group, which received usual care ([Black 2013](#); [Calland 2011](#); [Cheater 2005](#); [Curley 1998](#); [Deneckere 2013](#); [Schmidt 1998](#); [Strasser 2008](#); [Wild 2004](#)); one study compared two different IPC interventions ([Wilson 2004](#)).

All of the studies reported an objectively measured, or self-reported (using a validated instrument), patient, clinical process or efficiency outcome. The secondary outcome, collaborative behaviour, was only evaluated in four of the studies (see below).

Included studies

Externally facilitated interprofessional activities

[Black 2013](#) evaluated the effectiveness of an externally facilitated IPC intervention, based in primary care practices, to support nurses, administrative staff, practice managers, and receptionists to work collaboratively with general practitioners (GPs) to enhance the delivery of patient care when implementing practice systems that supported chronic disease care. Sixty primary care practices, involving 1185 participants based across Australia, took part in the study. The intervention included the following activities: structured appointment systems; patient disease registers; patient recall systems with reminders; patient education; planned care; definition of roles, responsibilities and job descriptions for each professional or staff member; communication and meetings; practice billing; record keeping; and quality improvement. Before the intervention, facilitators conducted workshops with primary care staff, followed by practice visits. Staff were provided with resource manuals and workbooks for each of the different elements of the intervention. Facilitators routinely followed up practices by telephone and email. All control teams received the IPC intervention after the 12-month follow-up data collection in each intervention practice.

[Cheater 2005](#) evaluated an externally facilitated programme aimed at improving IPC by the use of a multidisciplinary audit in a secondary care setting. Twenty-two multidisciplinary teams from five acute care hospitals in the UK participated. Each team consisted of nurses and physicians, and a representative from one or more of other health professional groups (e.g. pharmacist, social worker, physiotherapist), service support staff (e.g. ward clerk, care assistant), and managers (N = 141). A range of specialties were represented. After participating in a two-day skills workshop, external facilitators facilitated five meetings over a period of six months, for each of the multidisciplinary teams randomised to the intervention group. Intervention teams were required to undertake a

collaborative audit (the specific focus of these audits was not identified) and submit an audit report. Control teams provided usual care at their institution.

[Deneckere 2013](#) evaluated an interprofessional intervention for 30 teams caring for patients with chronic obstructive pulmonary disease and proximal femur fracture. Seventeen intervention teams and 13 control teams based in Belgian hospitals were involved in the study (N = 581), the aim of which was to examine effects of the use of care pathways on interprofessional teamwork, in an acute care setting. The intervention involved three facilitated components. To promote IPC the intervention included: feedback on the team's performance before the implementation of care pathways; receipt of evidence-based key indicators for implementing care pathways in practice; and review and training in the development and implementation of care pathways. The teams randomised to the control group provided usual care (i.e. did not implement care pathways). The intervention teams were given nine months to implement the intervention. After this time period, a summative evaluation was performed, in which the performance on team indicators was compared between the intervention and control groups.

[Strasser 2008](#) evaluated the effects of an externally facilitated intervention aimed at improving collaboration to support better care delivered to patients following a stroke. Thirty-one teams from different Veteran Affairs' rehabilitation units based in the USA participated in the study. Participants included physicians, nurses, occupational therapists, speech-language pathologists, physical therapists, case managers, or social workers (N = 464). Intervention teams received the following activities, delivered in three phases: phase 1) an off-site facilitated workshop emphasising team dynamics, problem-solving, the use of performance feedback data, and the development of action plans (specific team performance profiles with recommendations) for process improvement; phase 2) development of written action plans to address team process problems, based on discussions at the earlier workshop; and phase 3) telephone and video conference consultation on advice for implementation of action plans, and facilitation of team process skills. Control teams continued with usual care (i.e. specific team performance profile information).

For further details on these studies: see [Characteristics of included studies](#)

Interprofessional rounds

[Curley 1998](#) examined the effects of daily interdisciplinary rounds in inpatient medical wards at an acute care hospital in the USA. The intervention group consisted of three ward services that implemented interdisciplinary work rounds; the control group consisted of three other ward services that continued usual care (i.e.

traditional work rounds) (N = 1102). Team members included: medical interns and residents, staff nurses, nursing supervisors, respirologists, pharmacists, nutritionists, and social workers. To reduce baseline variability the authors used a process of random allocation of patient and clinical staff to either intervention wards or control wards.

[Wild 2004](#) studied the effects of daily interdisciplinary rounds in a telemetry unit of a community hospital in the USA. In this study, patients were randomised to a medical team that performed daily interdisciplinary rounds, and patients were randomised to a medical team that provided usual care (N = 84). During the interdisciplinary rounds, the resident physicians, nurses, a case manager, pharmacist, dietitian, and physical therapist spent two to five minutes discussing each patient, and identifying and addressing possible discharge problems. No information on the duration of the intervention was provided.

For further details on these studies: see [Characteristics of included studies](#).

Interprofessional meetings

[Schmidt 1998](#) evaluated the impact of multidisciplinary team meetings on the quality and quantity of psychotropic drug prescribing in Swedish nursing homes. Thirty-six nursing homes were randomised to either receive the intervention or were randomised to the control group (N = 1854). In the experimental nursing homes, the pharmacist in the homes helped organise team meetings that occurred approximately once a month, over a period of 12 months. The nursing home pharmacists attended two training sessions prior to, and three sessions during, the programme. The team meeting participants in the nursing homes included the pharmacist, a physician, and selected nurses and nursing assistants. All participants were encouraged to take part in the meeting discussions about the drug use of individual residents. Usual care continued to be used in the control homes. Nursing home residents' prescriptions were recorded one month before, and one month after, the 12-month intervention.

[Wilson 2004](#) compared two forms of multidisciplinary virtual team conferencing: usual care (audio conferencing) with video conferencing (including audio). Participating team members consisted of medical staff specialists, medical registrars, nurses, a

speech pathologist, occupational therapists, a social worker, and medical students. Patients were randomly assigned to the audio conferencing or video conferencing (N = 100). At each conference session, the audio conferences were conducted before the video conferences, with the same multidisciplinary team.

For further details on these studies: see [Characteristics of included studies](#).

Interprofessional checklists

[Calland 2011](#) studied the effects of a procedure checklist for interprofessional surgical teams during laparoscopic cholecystectomies. Ten USA-based general surgery teams were randomly assigned to an intervention or a control group (N = 29). The intervention consisted of preoperative steps: a briefing with introductions from all operating team members (surgeons, anaesthetists and nurses), a review of the patient's history, laboratory and radiographic studies, and a discussion of any unusual care circumstances. Surgeons in the intervention group received instructions on using the checklist and reminders before each surgery. In addition, a checklist copy was posted on the anaesthesia monitor in the operating room during cases, and team members were encouraged to use a call-and-repeat method to ensure that key steps of the checklist were neither omitted nor performed in a suboptimal manner. Control teams performed the laparoscopic cholecystectomy procedure in their normal fashion, without the use of a checklist.

For further details on this study: see [Characteristics of included studies](#).

Excluded studies

We excluded 30 studies. The main reason for exclusion was that the intervention was not practice-based interprofessional collaboration (IPC). See: [Characteristics of excluded studies](#).

Risk of bias in included studies

The risk of bias in the nine included studies varied. A brief summary is presented below. Further information for each included study is presented in the [Characteristics of included studies](#) tables; [Figure 2](#) and [Figure 3](#) provide further overviews.

Figure 2. 'Risk of bias' graph: review authors' judgements about each risk of bias item presented as percentages across all included studies, based on EPOC methods.




Figure 3. 'Risk of bias' summary: review authors' judgements about each risk of bias item for each included study, based on EPOC methods.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Baseline outcome measurements similar - All outcomes	Baseline characteristics similar	Blinding of outcome assessment (detection bias) - All outcomes	Incomplete outcome data (attrition bias) - All outcomes	Contamination	Selective reporting (reporting bias)	Other bias
Black 2013	?	?	+	+	-	-	+	?	+
Calland 2011	?	?	?	+	-	-	?	?	+
Cheater 2005	+	+	+	+	+	-	+	?	+
Curley 1998	+	+	?	+	+	+	+	?	+
Deneckere 2013	+	+	?	+	-	+	+	+	+
Schmidt 1998	?	+	+	+	+	+	+	?	+
Strasser 2008	+	?	+	+	-	-	+	+	+
Wild 2004	+	?	+	+	+	+	+	?	+
Wilson 2004	+	+	?	+	+	?	?	?	+

Allocation

We classified the sequence generation for six studies as 'low risk'. In [Cheater 2005](#) and [Strasser 2008](#), randomisation was determined by computer; in [Wild 2004](#), randomisation was performed with random numerical assignments in pre-sealed envelopes; [Curley 1998](#) randomised with a firm system; [Wilson 2004](#) used a table of random numbers, while [Deneckere 2013](#) used stratified randomisation to render intervention and control interprofessional teams comparable. We assessed the sequence generation for three studies as 'unclear risk', as there was insufficient information reported about the sequence generation process ([Black 2013](#); [Calland 2011](#); [Schmidt 1998](#)).

For allocation concealment, we classified five of the studies as 'low risk' ([Cheater 2005](#); [Curley 1998](#); [Deneckere 2013](#); [Schmidt 1998](#); [Wilson 2004](#)). In these studies, participating professionals or investigators enrolled participants. They could not foresee assignment because allocation took place in a location separate from recruitment; a sequentially numbered, blind or pre-sealed envelope was used for allocation concealment, or both. We classified four studies at 'unclear risk', since there was insufficient information provided to assess this criteria ([Black 2013](#); [Calland 2011](#); [Schmidt 1998](#); [Wild 2004](#)).

Blinding

We considered four studies to represent 'high risk' for blinding of outcome assessment ([Black 2013](#); [Calland 2011](#); [Deneckere 2013](#); [Strasser 2008](#)). These studies did not prevent knowledge of the allocated interventions, as no blinding was performed, and the outcome measurement was likely to be affected by lack of blinding. We classed detection bias as 'low risk' in five studies ([Cheater 2005](#); [Curley 1998](#); [Schmidt 1998](#); [Wild 2004](#); [Wilson 2004](#)).

Incomplete outcome data

We assessed four studies as being at 'low risk' of attrition bias ([Curley 1998](#); [Deneckere 2013](#); [Schmidt 1998](#); [Wild 2004](#)). These studies provided an adequate description of participant flow through the study, with missing outcome data relatively balanced between groups, and judged to be unlikely to be related to the outcomes of interest. We assessed four studies as having 'high risk' of attrition bias for the following reasons: they acknowledged that sites dropped out but intention-to-treat analysis was not mentioned in the text; there were differences in characteristics related to study outcomes between completers and non-completers ([Black 2013](#); [Calland 2011](#); [Cheater 2005](#); [Strasser 2008](#)). We judged attrition bias at 'unclear risk' in one study, as the study authors did not report sufficient information about attrition and missing data ([Wilson 2004](#)).

Selective reporting

Two of the more recently completed studies published research protocols prior to study commencement ([Deneckere 2013](#); [Strasser 2008](#)). We judged seven studies at 'unclear risk' for this bias, due to the difficulty in establishing what outcomes may have been planned in the protocol and collected, but not reported in the final published study report.

Other potential sources of bias

We assessed two of the studies as being at 'unclear risk' for contamination bias ([Calland 2011](#); [Wilson 2004](#)). In both trials, some measures were taken to prevent contamination, but some of the subjects crossed over from one arm of the study to the other (and were involved in both interventions), thereby, potentially contaminating the initial randomisation process. We recorded studies at 'unclear risk' for baseline outcome measurements and baseline characteristics due to lack of information presented in the published paper.

Effects of interventions

See: [Summary of findings for the main comparison](#) Effects of practice-based interprofessional collaboration (IPC) interventions on professional practice and healthcare outcomes compared to usual care; [Summary of findings 2](#) Effects of practice-based interprofessional collaboration (IPC) interventions on professional practice and healthcare outcomes compared with alternative IPC intervention

This section reports on primary and secondary outcomes from the nine included studies. Data presented below are taken directly from the published articles of the included studies. We report point estimates and confidence intervals whenever reported by the study authors. For further detailed information see [Summary of findings for the main comparison](#) and [Summary of findings 2](#).

Practice-based interprofessional collaboration (IPC) interventions compared with usual care

Eight studies compared IPC interventions with usual care.

Externally facilitated interprofessional activities

Patient health outcomes

Patient functional status

Externally facilitated interprofessional activities may slightly improve stroke patients' motor function (low-certainty evidence, 1 study, N = 464). [Strasser 2008](#) reported a difference between the intervention and control groups in patient health, with changes from admission to discharge in the motor skills component of the Functional Independence Measure (FIM) score. For the patients with stroke, 13.6% more of those in the intervention group gained in excess of the median gain, 23 points, when compared to the control group (P = 0.032).

Patient-assessed quality of care

It is uncertain if externally facilitated interprofessional activities increase patient-assessed quality of care at 12-month follow-up (very low-certainty evidence, 1 study, N = 1185) ([Black 2013](#)).

Patient mortality, morbidity or complication rates

None of the included studies reported patient mortality, morbidity or complication rates.

Clinical process or efficiency outcomes

Adherence to recommended practices

The use of interprofessional activities with an external facilitator may slightly improve adherence to recommended practices and prescription of drugs (low-certainty evidence, 2 studies, N = 722). [Cheater 2005](#) reported an increase in collaborative audit activity, with six of the 11 intervention teams completing the full audit cycle. Only three control teams undertook any audit (first data collection).

[Deneckere 2013](#) reported improvements in the following outcomes: conflict management (slope of difference between intervention and control group (β) 0.30, 95% confidence interval (CI) 0.08 to 0.53); team climate for innovation (β 0.29, 95% CI 0.09 to 0.49); and level of organised care (β 5.56, 95% CI 1.35 to 9.76). [Deneckere 2013](#) also reported that the intervention group scored lower in emotional exhaustion (β 0.57, 95% CI 0.14 to 1.00) and higher in level of competence (β 0.39, 95% CI 0.15 to 0.64).

Continuity of care

It is uncertain if externally facilitated interprofessional activities improve continuity of care (low-certainty evidence, 1 study, N = 464).

Use of healthcare resources

While [Strasser 2008](#) reported that externally facilitated interprofessional activities may improve use of resources (low-certainty evidence, 1 study, N = 464), [Strasser 2008](#) also reported that there was little or no difference between the intervention and control groups for length of stay or discharge disposition.

Collaborative behaviour outcomes

Collaborative working, team communication and team co-ordination

It is uncertain whether externally facilitated interprofessional activities improve collaborative working, team communication, and co-ordination (very low-certainty evidence, 3 studies, N = 1907). [Black 2013](#) reported differences between the intervention and control groups in the mean change from baseline to follow-up in staff role scores assessed using the Chronic Care Team Profile (CCTP). These differences were in the non-GP clinical staff function (P = 0.023), the administrative staff function (P < 0.001), and the total score (P = 0.03). These changes included, for example, the creation of a diabetes care co-ordinator to perform tasks such as managing the recall and reminder system for patients with diabetes, and organising staff meetings to improve communication and practice systems.

[Cheater 2005](#) report that Collaborative Practice Scale (CPS) scores on co-operation went from 83.5 at baseline to 88.5 after the intervention in the intervention group, compared to 84.5 at baseline to 84.5 after the intervention in the control group. These differences are presented with no measures of variability and the change appears small in relation to the value at baseline.

[Deneckere 2013](#) found little effect of the intervention on relational co-ordination, which assessed the process of communication and relationship between team members in order to complete the task.

Interprofessional rounds

Patient health outcomes

None of the included studies reported patient health outcomes.

Clinical process or efficiency outcomes

Use of healthcare resources

Interprofessional rounds may improve use of healthcare resources (low-certainty evidence, 2 studies, N = 1186).

[Curley 1998](#) found differences in length of stay and costs for patients in the interdisciplinary group compared to the traditional

care group. The mean length of stay for patients in the interdisciplinary rounds group was 5.46 days, compared with 6.06 days for traditional care ($P = 0.006$). The mean total charges were USD 6681 and USD 8090 for the two groups, respectively ($P = 0.002$). For respiratory therapy, 91.7% of the nursing and pharmacy orders in the interdisciplinary rounds group were appropriate, compared with 73.6% for the traditional rounds group ($P = 0.075$). [Wild 2004](#) found little or no change in the length of hospital stay, between the experimental group (3.2 ± 2.7 days), which participated in interdisciplinary rounds, and the control group (3.2 ± 3.2 days ($P = 0.90$)).

Collaborative behaviour outcomes

None of the included studies reported collaborative behaviour outcomes.

Interprofessional meetings

Patient health outcomes

The included study did not report patient health outcomes.

Clinical process or efficiency outcomes

Adherence to recommended practices

Interprofessional meetings may slightly improve adherence to recommended practices and prescription of drugs (low-certainty evidence, 1 study, $N = 1854$).

[Schmidt 1998](#) found that the change from baseline to end of study in the proportion of patients receiving drugs was the same in experimental and control homes (1.3% intervention and 1.3% control). The mean number of psychotropic drugs increased from 2.07 to 2.08 (1%) in the intervention group and from 2.06 to 2.20 (7%) in the control group. The use of non-recommended hypnotics declined by 37% in the experimental homes versus a decrease of only 3% in the control homes. There was little or no change in the prescribing of non-recommended anxiolytics in the experimental homes, but there was an increase of 7% in the control homes. Non-recommended antidepressant drugs decreased by 59% in experimental homes but by only 34% in control homes.

Collaborative behaviour outcomes

The included study did not report collaborative behaviour outcomes.

Interprofessional checklists

Patient health outcomes

None of the included studies reported patient health outcomes.

Clinical process or efficiency outcomes

Use of healthcare resources

Interprofessional checklists may improve use of healthcare resources (low-certainty evidence, 1 study, $N = 29$).

In [Calland 2011](#), there was little or no difference between surgeons and team members in the intervention group (who received basic team training and used a pre-procedural checklist), and the control group (who performed standard laparoscopic cholecystectomies), in patient outcomes, length of operation, discharge status, readmission rates, and technical proficiency. Overall, situational awareness did not differ between the two groups. Surgeons and team members in the intervention group consistently rated their cases as involving less satisfactory levels of comfort, team efficiency, and communication compared to the control group.

Collaborative behaviour outcomes

Collaborative working, team communication and team coordination

[Calland 2011](#) reported that participants in the intervention (checklist) group were more likely to introduce team members, assign team roles, give case presentations, devise contingency plans, and complete post-case performance reviews. The intervention may make little or no difference to collaborative behaviour.

Practice-based IPC intervention compared with alternative IPC intervention

[Wilson 2004](#) compared one type of IPC intervention (video conferencing) with a second IPC intervention (audio conferencing).

Interprofessional meetings

Patient health outcomes

The included study did not report patient health outcomes.

Clinical process or efficiency outcomes

Use of healthcare resources

Interprofessional meetings may improve use of healthcare resources (low-certainty evidence, 1 study, N = 100). [Wilson 2004](#) reported that the mean number of audio conferences held per patient (mean 3.3; standard deviation (SD) 4.4) was greater than the mean number of video conferences held (mean 1.9; SD 1.3; P = 0.04). Video conferencing may reduce the average length of treatment (mean 6.0; SD 4.5 days), compared to audio conferencing (mean 10.2; SD 12.3 days; P = 0.03). The use of video confer-

encing may improve process/efficiency outcomes by reducing the number of multidisciplinary conferences needed per patient and patient length of stay.

Collaborative behaviour outcomes

Collaborative working, team communication and team coordination

[Wilson 2004](#) reported little or no difference between the groups in the number of communications between health professionals, as recorded in the notes (low-certainty evidence, 1 study, N = 100).

ADDITIONAL SUMMARY OF FINDINGS *[Explanation]*

Effects of practice-based interprofessional collaboration (IPC) interventions on professional practice and healthcare outcomes compared with alternative IPC intervention

Patient or population: health and social care professionals involved in the delivery of health services and patient care

Settings: two hospitals in Australia

Intervention: multidisciplinary video conferencing

Comparison: multidisciplinary audio conferencing

Outcomes	Impacts	No. of studies (participants)	Certainty of the evidence (GRADE)
Patient health outcomes	The study did not report patient health outcomes.	-	-
Clinical process or efficiency outcomes	Video conferencing may reduce the average length of treatment, compared to audio conferencing and may improve process/efficiency outcomes by reducing the number of multidisciplinary conferences needed per patient and patient length of stay	1 (100)	⊕⊕⊕⊕ Low ^a
Collaborative behaviour outcomes	There was little or no difference between the interventions in the number of communications between health professionals	1 (100)	⊕⊕⊕⊕ Low ^a

GRADE Working Group grades of evidence

High-certainty: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate-certainty: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low-certainty: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low-certainty: We are very uncertain about the estimate.

^a We assessed the certainty of evidence as low because of high risk of bias (attrition and detection) and unclear risk of bias (selection, reporting, and contamination).

Summary of main results

Eight studies compared an interprofessional collaboration (IPC) intervention with usual care ([Black 2013](#); [Calland 2011](#); [Cheater](#)

DISCUSSION

2005; Curley 1998; Deneckere 2013; Schmidt 1998; Strasser 2008; Wild 2004). One study compared one IPC intervention (video/audio conferencing) with another (audio conferencing) (Wilson 2004). See [Summary of findings for the main comparison](#) and [Summary of findings 2](#).

IPC interventions compared with usual care

Each of the eight included studies in this comparison is discussed below in relation to the specific IPC intervention they employed. None of the included studies reported on patient mortality, morbidity or complication rates.

Externally facilitated interprofessional activities

Four studies reported the effects of this intervention. Externally facilitated interprofessional activities may slightly improve patient functional status (1 study, N = 464) and adherence to recommended practices (2 studies, N = 722), and may improve use of resources (1 study, N = 464). There was low-certainty evidence for all outcomes. It is uncertain whether externally facilitated interprofessional activities improve patient-assessed quality of care (1 study, N = 1185), continuity of care (1 study, N = 464), or collaborative working (3 studies, N = 1907), as we graded the evidence as very low-certainty.

The included studies reported varied activities. Strasser 2008 implemented a multiphase, IPC programme, delivered over six months, that aimed to enhance the effectiveness of interdisciplinary stroke rehabilitation teams. Cheater 2005 reported an IPC intervention where an external facilitator used strategies to encourage collaborative working. Deneckere 2013 developed a care pathway for patients hospitalised in an acute hospital setting with either a proximal femur fracture or an exacerbation of chronic obstructive pulmonary disease. Black 2013 used a multimodal intervention involving educational sessions, practice visits, and resource manuals and workbooks.

Interprofessional rounds

Two studies reported the effects of this intervention. Overall, interprofessional rounds may improve use of resources (low-certainty evidence, 2 studies, N = 1186 participants). The studies implemented the intervention in an acute care hospital and in a community hospital telemetry ward (Curley 1998; Wild 2004). Wild 2004 suggested that their finding of little or no change in clinical process outcomes could be because, for many admissions, there was already a clinical pathway with standardised care for their diagnoses, the patients were more stable, at a lower risk for complications, and possibly healthier overall, and so the interdisciplinary rounds provided no additional advantage.

Interprofessional meetings

One study reported the effects of this intervention. Interprofessional meetings may slightly improve adherence to recommended practices (low-certainty evidence, 1 study, N = 1854) and may improve use of resources (low-certainty evidence, 1 study). Schmidt 1998 implemented a collaborative team meeting in nursing homes.

Interprofessional checklists

One study reported the effects of this intervention. Interprofessional checklists may improve the use of resources (low-certainty evidence, 1 study, N = 29). Calland 2011 used a pre-procedural checklist with surgical teams.

IPC intervention compared with alternative IPC intervention

Wilson 2004 assessed the impacts of interprofessional meetings facilitated using two different technologies, and found that video/audio conferencing may be more efficient than audio conferencing alone.

Overall completeness and applicability of evidence

The included studies covered four types of practice-based IPC interventions: externally facilitated interprofessional activities, interprofessional rounds, interprofessional meetings, and interprofessional checklists. We did not find any studies that used other types of practice-based IPC interventions, such as debriefing. Given the range of practice-based interventions aimed at promoting IPC, and the different types of participants, settings, and clinical areas addressed in these interventions, further studies are required to provide better insight into the effectiveness of these interventions alone or in combination, in a variety of target groups and clinical areas.

Other issues affected the completeness of the evidence. For example, Schmidt 1998 acknowledged that their study could not provide robust evidence about the participating teams' decision-making processes, or the strategies used by pharmacists in their role as team facilitators.

Secondary outcomes, focused on examining interprofessional collaboration processes, were not well examined in most of the studies, with only four studies reporting on this type of outcome (Black 2013; Cheater 2005; Deneckere 2013; Wilson 2004).

Six of the nine included studies were cluster-randomised trials; this was appropriate, given the complex nature of interventions and their inherently clustered nature, the difficulty of blinding, and the consequential threat of contamination.

Whilst we identified four new studies for this review, the number of practice-based IPC studies remains small. Some of the studies offered some evidence that IPC interventions may be effective in

improving clinical processes/efficiency outcomes, but the small number of studies and the methodological limitations precluded definitive conclusions. Therefore, we still know little about the processes of collaboration, and how they contribute to changes in clinical process/efficiency and patient outcomes.

Certainty of the evidence

The review included nine randomised trials; the findings for the two comparisons are summarised in [Summary of findings for the main comparison](#) and [Summary of findings 2](#). Based on our GRADE assessment, we found the certainty of the evidence from the included studies to be low or very low due to risk of bias (attrition, detection, selection, reporting, and contamination bias), and potential indirectness (as outcomes generated in one country or clinical setting may not be transferable to other settings).

A number of other study limitations may also have contributed to the risk of bias: [Curley 1998](#) used a non-validated survey to examine interdisciplinary communication on the ward. [Wild 2004](#) used a questionnaire to ask about communication, but this was only administered to the experimental group. Similarly, [Cheater 2005](#) used a modified Collaborative Practice Scale (CPS), which was also only completed by the experimental group. [Wilson 2004](#) used the number of communications between health professionals, recorded in the notes, to measure communication, which is a limited measurement of collaboration. [Deneckere 2013](#) used a post-test only; as a result, there was no baseline assessment measured before the intervention. Also, the results were primarily based on self-reported outcomes, which caused some limitations, such as possible social desirability bias.

Potential biases in the review process

The searches were sensitive, but some literature may be under-represented. We did not contact authors of the included studies for this review, which may have introduced some bias. In the next review, Scopus should be added to the databases to be searched, and corresponding authors of included studies should be contacted to clarify published information, and to seek unpublished data. The limited number of studies reporting data on both primary and secondary outcomes limited exploration of publication bias and sensitivity analyses.

Agreements and disagreements with other studies or reviews

There were no comparable reviews in this area.

AUTHORS' CONCLUSIONS

Implications for practice

The findings from the nine studies included in this review suggested that interventions aimed at improving interprofessional collaboration (IPC) through practice changes may slightly improve clinical process/efficiency and patient health outcomes compared to usual care or an alternative intervention. Nine randomised trials of four different IPC interventions, in nine different clinical settings and health conditions provided mixed results. We judged the certainty of evidence from these randomised trials to be low to very low. Based on these included studies we do not have sufficient evidence to draw clear conclusions on the effects of IPC interventions.

Implications for research

Given the problems that health professionals encounter with IPC in their clinical practice (e.g. [Körner 2016](#); [Van Leijen-Zeelenberg 2015](#)), it is encouraging that research on the effects of IPC interventions has increased since the previous Cochrane Review of this intervention ([Zwarenstein 2009](#)). While this research field is developing, further rigorous, mixed-method studies are required. It is recommended that future randomised trials have a clear and explicit focus on IPC, longer acclimatisation periods before evaluating newly implemented teamwork interventions, and longer follow-up.

Future research should also focus on the conceptualisation and measurement of collaboration. While there are some scales that measure collaboration (e.g. [Kenaszchuk 2010](#)), there are limitations with their validity, reliability, the extent to which they could be used with different professional groups, and how well they examine issues of collaborative practice.

The studies included in this updated Cochrane Review used a variety of terms to describe their interventions (e.g. interdisciplinary, multidisciplinary), which contributes to an on-going confusion of terminology. The absence of a consistent approach to terminology of these interventions undermines our ability to synthesise them in order to develop a more informed understanding of their effects. Further work is needed to clarify the conceptualisation of IPC, interprofessional education, and interprofessional organisationally-based interventions to support consistency in the use and understanding of these terms and their related interventions. While we have published an initial classification ([Reeves 2010](#); [Reeves 2011](#)), future empirical work could test these conceptualisations to generate more detailed knowledge related to their implementation. Finally, quantitative and qualitative methods should be used in single studies to improve our understanding of how the intervention addresses collaboration, the nature of changes that occur in relation to collaboration, and how they in turn lead to the outcomes achieved.

ACKNOWLEDGEMENTS

The authors would like to acknowledge their respective academic institutions for the support in undertaking this updated Cochrane Review. The authors would also like to thank the following Effective Practice and Organisation of Care (EPOC) editors and staff for their advice and feedback on the review: Pierre Durieux, Daniela Gonçalves Bradley, Simon Lewin, Paul Miller, and Julia Worswick. We would like to thank Vicki Pennick for copy-editing the review. Finally, the authors would like to thank the peer re-

viewers Kunal D Patel and Andreas Xyrichis who generously gave up their time to provide comments on the review, which helped improve its quality.

The National Institute for Health Research (NIHR), via Cochrane Infrastructure, funds the Effective Practice and Organisation of Care (EPOC) Group. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, National Health Service (NHS), or the Department of Health.

REFERENCES

References to studies included in this review

Black 2013 *{published data only}*

* Black DA, Taggart J, Jayasinghe UW, Proudfoot J, Crookes P, Beilby J, et al. Teamwork Research Team. The Teamwork Study: enhancing the role of non-GP staff in chronic disease management in general practice. *Australian Journal of Primary Health* 2013;**19**(3):184–9.

Calland 2011 *{published data only}*

* Calland JF, Turrentine FE, Guerlain S, Bovbjerg V, Poole GR, Lebeau K, et al. The surgical safety checklist: lessons learned during implementation. *The American Surgeon* 2011;**77**(9):1131–7.

Cheater 2005 *{published data only}*

Cheater FM, Hearnshaw H, Baker R, Keane M. Can a facilitated programme promote effective multidisciplinary audit in secondary care teams? An exploratory trial. *International Journal of Nursing Studies* 2005;**42**:779–91.

Curley 1998 *{published data only}*

* Curley C, McEachern JE, Speroff T. A firm trial of interdisciplinary rounds on the inpatient medical wards. *Medical Care* 1998;**36**(8 Suppl):AS4–12.

Deneckere 2013 *{published data only}*

Deneckere S, Euwema M, Lodewijckx C, Panella M, Mutsvari T, Sermeus W, et al. Better interprofessional teamwork, higher level of organized care, and lower risk of burnout in acute health care teams using care pathways: a cluster randomized controlled trial. *Medical Care* 2013;**51**(1):99–107.

Schmidt 1998 *{published data only}*

Schmidt I, Claesson CB, Westerholm B, Nilsson LG, Svarstad BL. The impact of regular multidisciplinary team interventions on psychotropic prescribing in Swedish nursing homes. *Journal of the American Geriatrics Society* 1998;**46**:77–82.

Strasser 2008 *{published data only}*

Strasser DC, Falconer JA, Stevens AB, Uomoto JM, Herrin J, Bowen SE, et al. Team training and stroke rehabilitation outcomes: a cluster randomized trial. *Archives of Physical Medicine and Rehabilitation* 2008;**89**(1):10–5.

Wild 2004 *{published data only}*

Wild D, Nawaz H, Chan W, Katz DL. Effects of interdisciplinary rounds on length of stay in a telemetry unit. *Journal of Public Health Management and Practice* 2004;**10**:63–9.

Wilson 2004 *{published data only}*

Wilson SF, Marks R, Collins N, Warner B, Frick L. Benefits of multidisciplinary case conferencing using audiovisual compared with telephone communication: a randomized controlled trial. *Journal of Telemedicine and Telecare* 2004;**10**:351–4.

References to studies excluded from this review

Bekelman 2015 *{published data only}*

Bekelman DB, Plomondon ME, Carey EP, Sullivan MD, Nelson KM, Hattler B, et al. Primary results of the patient-centered disease management (PCDM) for heart failure study: a randomized clinical trial. *JAMA Internal Medicine* 2015;**175**(5):725–32.

Boet 2013 *{published data only}*

* Boet S, Bould MD, Sharma B, Reeves S, Naik VN, Tribby E, et al. Within-team debriefing versus instructor-led debriefing for simulation-based education: a randomized controlled trial. *Annals of Surgery* 2013;**258**(1):53–8.

Boone 2008 *{published data only}*

* Boone BN, King ML, Gresham LS, Wahl P, Suh E. Conflict management training and nurse-physician collaborative behaviors. *Journal for Nurses in Staff Development* 2008;**24**(4):168–75.

Chen 2013 *{published data only}*

Chen Z, Ernst ME, Ardery G, Xu Y, Carter BL. Physician-pharmacist co-management and 24-hour blood pressure control. *Journal of Clinical Hypertension* 2013;**15**(5):337–443.

Cheng 2013 *{published data only}*

* Cheng A, Hunt EA, Donoghue A, Nelson-McMillan K, Nishisaki A, Leflore J, et al. EXPRESS Investigators. Examining pediatric resuscitation education using simulation and scripted debriefing: a multicenter randomized trial. *JAMA Pediatrics* 2013;**167**(6):528–36.

- Curtis 2012** *{published data only}*
Curtis JR, Ciechanowski PS, Downey L, Gold J, Nielsen EL, Shannon SE, et al. Development and evaluation of an interprofessional communication intervention to improve family outcomes in the ICU. *Contemporary Clinical Trials* 2012;**33**(6):1245–54.
- Dhalla 2014** *{published data only}*
Dhalla IA, O'Brien T, Morra D, Thorpe KE, Wong BM, Mehta R, et al. Effect of a post-discharge virtual ward on readmission or death for high-risk patients: a randomized clinical trial. *JAMA* 2014;**312**(13):1305–12.
- Döpp 2013** *{published data only}*
Döpp CM, Graff MJ, Teerenstra S, Nijhuis-van der Sanden MW, Olde Rikkert MG, Vernooij-Dassen MJ. Effectiveness of a multifaceted implementation strategy on physicians' referral behavior to an evidence-based psychosocial intervention in dementia: a cluster randomized controlled trial. *BMC Family Practice* 2013;**14**:70. [DOI: 10.1186/1471-2296-14-70]
- Fransen 2012** *{published data only}*
Fransen AF, van de Ven J, Meriën AE, de Wit-Zuurendonk LD, Houterman S, Mol BW, et al. Effect of obstetric team training on team performance and medical technical skills: a randomised controlled trial. *BJOG* 2012;**119**(11):1387–93.
- Goud 2009** *{published data only}*
Goud R, de Keizer NF, ter Riet G, Wyatt JC, Hasman A, Hellemans IM, et al. Effect of guideline based computerised decision support on decision making of multidisciplinary teams: cluster randomised trial in cardiac rehabilitation. *BMJ* 2009;**338**:b1440. [DOI: 10.1136/bmj.b1440]
- Hallin 2011** *{published data only}*
Hallin K, Henriksson P, Dalén N, Kiessling A. Effects of interprofessional education on patient perceived quality of care. *Medical Teacher* 2011;**33**(1):e22–6.
- Hobgood 2010** *{published data only}*
Hobgood C, Sherwood G, Frush K, Hollar D, Maynard L, Foster B, et al. Interprofessional Patient Safety Education Collaborative. Teamwork training with nursing and medical students: does the method matter? Results of an interinstitutional, interdisciplinary collaboration. *Quality & Safety in Health Care* 2010;**19**(6):e25.
- Hoffmann 2014** *{published data only}*
Hoffmann B, Müller V, Rochon J, Gondan M, Müller B, Albay Z, et al. Effects of a team-based assessment and intervention on patient safety culture in general practice: an open randomised controlled trial. *BMJ Quality & Safety* 2014;**23**(1):35–46.
- Jankouskas 2011** *{published data only}*
* Jankouskas TS, Haidet KK, Hupcey JE, Kolanowski A, Murray WB. Targeted crisis resource management training improves performance among randomized nursing and medical students. *Simulation in Healthcare* 2011;**6**(6): 316–26.
- Jenkins 2013** *{published data only}*
Jenkins VA, Farewell D, Farewell V, Batt L, Wagstaff J, Langridge C, et al. Teams Talking Trials Steering Committee. Teams Talking Trials: results of an RCT to improve the communication of cancer teams about treatment trials. *Contemporary Clinical Trials* 2013;**35**(1): 43–51.
- Katakam 2012** *{published data only}*
Katakam LI, Trickey AW, Thomas EJ. Speaking up and sharing information improves trainee neonatal resuscitations. *Journal of Patient Safety* 2012;**8**(4):202–9.
- Keller 2012** *{published data only}*
Keller C, Records K, Coe K, Ainsworth B, Vega López S, Nagle-Williams A, et al. Promotoras' roles in integrative validity and treatment fidelity efforts in randomized controlled trials. *Family & Community Health* 2012;**35**(2): 120–9.
- Kemper 2011** *{published data only}*
Kemper PF, de Bruijne M, Van Dyck C, Wagner C. Effectiveness of classroom based crew resource management training in the intensive care unit: study design of a controlled trial. *BMC Health Services Research* 2011;**11**:304.
- Koerner 2014** *{published data only}*
Koerner M, Wirtz M, Michaelis M, Ehrhardt H, Steger AK, Zerpies E, et al. A multicentre cluster-randomized controlled study to evaluate a train-the-trainer programme for implementing internal and external participation in medical rehabilitation. *Clinical Rehabilitation* 2014;**28**(1): 20–35.
- Körner 2012** *{published data only}*
Körner M, Ehrhardt H, Steger AK, Bengel J. Interprofessional SDM train-the-trainer program “Fit for SDM”: provider satisfaction and impact on participation. *Patient Education and Counseling* 2012;**89**(1):122–8.
- Kunkler 2007** *{published data only}*
Kunkler IH, Prescott RJ, Lee RJ, Brebner JA, Cairns JA, Fielding RG, et al. TELEMAM: a cluster randomised trial to assess the use of telemedicine in multi-disciplinary breast cancer decision making. *European Journal of Cancer* 2007;**43**(17):2506–14.
- Lee 2013** *{published data only}*
* Lee GA, Murray A, Bushnell R, Niggemeyer LE. Challenges developing evidence-based algorithms for the trauma reception and resuscitation project. *International Emergency Nursing* 2013;**21**(2):129–35.
- Marsteller 2013** *{published data only}*
Marsteller JA, Hsu YJ, Wen M, Wolff J, Frick K, Reider L, et al. Effects of Guided Care on providers' satisfaction with care: a three-year matched-pair cluster-randomized trial. *Population Health Management* 2013;**16**(5):317–25.
- Mohaupt 2012** *{published data only}*
* Mohaupt J, van Soeren M, Andrusyszyn MA, Macmillan K, Devlin-Cop S, Reeves S. Understanding interprofessional relationships by the use of contact theory. *Journal of Interprofessional Care* 2012;**26**(5):370–5.
- Musick 2011** *{published data only}*
Musick BS, Robb SL, Burns DS, Stegenga K, Yan M, McCorkle KJ, et al. Development and use of a web-based

data management system for a randomized clinical trial of adolescents and young adults. *Computers, Informatics, Nursing: CIN* 2011;**29**(6):337–43.

O’Leary 2010 *{published data only}*

O’Leary KJ, Wayne DB, Haviley C, Slade ME, Lee J, Williams MV. Improving teamwork: impact of structured interdisciplinary rounds on a medical teaching unit. *Journal of General Internal Medicine* 2010;**25**(8):826–32.

Rörtgen 2013 *{published data only}*

* Rörtgen D, Bergrath S, Rossaint R, Beckers SK, Fischermann H, Na IS, et al. Comparison of physician staffed emergency teams with paramedic teams assisted by telemedicine – a randomized, controlled simulation study. *Resuscitation* 2013;**84**(1):85–92.

Van de Ven 2010 *{published data only}*

* Van de Ven J, Houterman S, Steinweg RA, Scherpbier AJ, Wijers W, Mol BW, et al. TOSTI-Trial Group. Reducing errors in health care: cost-effectiveness of multidisciplinary team training in obstetric emergencies (TOSTI study); a randomised controlled trial. *BMC Pregnancy Childbirth* 2010;**10**:59.

Weller 2014 *{published data only}*

* Weller JM, Torrie J, Boyd M, Frengley R, Garden A, Ng WL, et al. Improving team information sharing with a structured call-out in anaesthetic emergencies: a randomized controlled trial. *British Journal of Anaesthesia* 2014;**112**(6):1042–9.

Wittenberg-Lyles 2013 *{published data only}*

* Wittenberg-Lyles E, Oliver DP, Kruse RL, Demiris G, Gage LA, Wagner K. Family caregiver participation in hospice interdisciplinary team meetings: how does it affect the nature and content of communication?. *Health Communication* 2013;**28**(2):110–8.

Additional references

Baker 2011

Baker L, Egan-Lee E, Martimianakis M, Reeves S. Relationships of power: implications for interprofessional education and practice. *Journal of Interprofessional Care* 2011;**25**:98–104.

Department of Health 1997

UK Department of Health. The New NHS: Modern and Dependable. Available from www.gov.uk/government/publications/the-new-nhs.

EPOC 2013

Cochrane Effective Practice, Organisation of Care (EPOC). EPOC worksheets for preparing a ‘Summary of findings’ table using GRADE. EPOC resources for review authors, 2013. Available from epoc.cochrane.org/epoc-specific-resources-review-authors.

EPOC 2016

Cochrane Effective Practice, Organisation of Care (EPOC). Suggested risk of bias criteria for EPOC reviews. EPOC resources for review authors, 2016. Available from epoc.cochrane.org/epoc-specific-resources-review-authors.

Francis 2013

Francis R. Report of the Mid Staffordshire NHS Foundation Trust public inquiry. Available from www.gov.uk/government/uploads/system/uploads/attachment_data/file/279124/0947.pdf.

Guyatt 2008

Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schünemann HJ, GRADE Working Group. What is “quality of evidence” and why is it important to clinicians?. *BMJ* 2008;**336**(7651):995–8.

Health Canada 2003

Health Canada. 2003 First Ministers’ Accord on Health Care Renewal. Available from www.scics.gc.ca/CMFiles/800039004_e1GTC-352011-6102.pdf.

Higgins 2011

Higgins JP, Green S, editor(s). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Institute of Medicine 2000

Institute of Medicine (US) Committee on Quality of Health Care in America, Kohn LT, Corrigan JM, Donaldson MS, editor(s). *To Err Is Human: Building a Safer Health System*. Washington (DC): National Academies Press (US), 2000.

Institute of Medicine 2013

Global Forum on Innovation in Health Professional Education, Board on Global Health, Institute of Medicine. *Interprofessional Education for Collaboration: Learning how to improve health from interprofessional models across the continuum of education to practice: Workshop summary*. Washington (DC): The National Academies Press (US), 2013.

Kenaszchuk 2010

Kenaszchuk C, Reeves S, Nicholas D, Zwarenstein M. Validity and reliability of a multiple-group measurement scale for interprofessional collaboration. *BMC Health Services Research* 2010;**10**:83. [DOI: 10.1186/1472-6963-10-83]

Körner 2016

Körner M, Bütof S, Müller C, Zimmermann L, Becker S, Bengel J. Interprofessional teamwork and team interventions in chronic care: A systematic review. *Journal of Interprofessional Care* 2016;**30**(1):15–28.

Lillebo 2015

Lillebo B, Faxvaag A. Continuous interprofessional coordination in perioperative work: an exploratory study. *Journal of Interprofessional Care* 2015;**29**(2):125–30.

Reeves 2010

Reeves S, Lewin S, Espin S, Zwarenstein M. *Interprofessional Teamwork for Health and Social Care*. London: Blackwell-Wiley, 2010.

Reeves 2011

Reeves S, Goldman J, Gilbert J, Tepper J, Silver I, Suter E, et al. A scoping review to improve conceptual clarity of

interprofessional interventions. *Journal of Interprofessional Care* 2011;**25**:167–74.

Reeves 2013

Reeves S, Perrier L, Goldman J, Freeth D, Zwarenstein M. Interprofessional education: effects on professional practice and healthcare outcomes (update). *Cochrane Database of Systematic Reviews* 2013, Issue 3. [DOI: 10.1002/14651858.CD002213.pub3]

The Joint Commission 2016

The Joint Commission (USA). Sentinel Event Data - Root Causes by Event Type. www.jointcommission.org/se/data/event/type/by/year/.

Van Leijen-Zeelenberg 2015

Van Leijen-Zeelenberg J, Van Raak A, Duimel-Peters I, Kroese M, Brink P, Vrijhoef H. Interprofessional communication failures in acute care chains: How can we identify the causes?. *Journal of Interprofessional Care* 2015; **29**(4):320–30.

WHO 1976

World Health Organization. Continuing Education of Health Personnel, 1976. www.who.int/genomics/professionals/education/en/.

WHO 2010

World Health Organization. Framework for action on interprofessional education and collaborative practice, 2010. www.who.int/hrh/resources/framework-action/en/.

References to other published versions of this review

Zwarenstein 1996

Zwarenstein M, Bryant W, Baillie R. The effects on patient care of interventions to change collaboration between nurses and doctors. *Cochrane Database of Systematic Reviews* 1996, Issue 3. [DOI: 10.1002/14651858.CD000072]

Zwarenstein 1997

Zwarenstein M, Bryant W, Baillie R, Sibthorpe B. Interventions to change collaboration between nurses and doctors. *Cochrane Database of Systematic Reviews* 1997, Issue 2. [DOI: 10.1002/14651858.CD000072.pub2]

Zwarenstein 2009

Zwarenstein M, Goldman J, Reeves S. Interprofessional collaboration: effects of practice-based interventions on professional practice and healthcare outcomes. *Cochrane Database of Systematic Reviews* 2009, Issue 8. [DOI: 10.1002/14651858.CD000072.pub2]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies *[ordered by study ID]*

Black 2013

Methods	Cluster-randomised trial to test the effectiveness of an intervention involving non GP-staff in GP practices, on the quality of care for patients with diabetes or cardiovascular disease
Participants	Country: Australia General practitioners, nurses, practice managers, receptionists, and other administrative staff. 60 general practices were randomised to receive a 6-month teamwork intervention immediately (intervention, n = 637) or after 12 months (control, n = 548)
Interventions	To assist non-GP staff (e.g. nurses, administrative staff (practice managers, receptionists)) to work as a team with GPs, the intervention included a number of activities including: the use of structured appointment systems, recall and reminders, planned care, the use of roles, responsibilities, and job descriptions, as well as communication and meetings
Outcomes	Quality of care (12-month follow-up)
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomisation is mentioned: "...Following baseline-data collection, practices were stratified according to size (solo, 2 to 4 GPs or 5+ GPs) and randomised to receive the 6-month teamwork intervention immediately, or after 12 months...", but method not specified
Allocation concealment (selection bias)	Unclear risk	No description of allocation method.
Baseline outcome measurements similar - All outcomes	Low risk	At baseline, the quality of care PACIC outcomes in the intervention group (3.01, SD 0.30) and control group (2.87, SD 0.34) were similar
Baseline characteristics similar	Low risk	Intervention and control teams look reasonably similar. Quote: "Control practices were more likely to be in an urban location compared with the intervention practices, have a lower full-time equivalent level of practice nurses and were also more likely to have a higher score

		on the CCTP with more administrative functions for chronic disease managed by non-GP staff. There were no key differences between the control and intervention practices for total levels of non-GP staffing.”
Blinding of outcome assessment (detection bias) - All outcomes	High risk	It did not appear that there was any blinding.
Incomplete outcome data (attrition bias) - All outcomes	High risk	Acknowledged sites dropped out, but ITT is not mentioned in the text Practice level: Quote: “Of these, 69% (60/87) finally participated in the study, and three of these (3/60) withdrew at follow up...Reasons for withdrawal of three practices included concern about the extent of data collection and other reasons not pertaining to the study.” Patient level: There were 3349 patients invited to participate in the study, with 2642 (79%) providing informed consent. Of these, 2552 (96.6%) returned the PACIC questionnaire at baseline, with 2135 (73.7%) completing all 20 items. To be included in the factor analysis, at least 17 questions needed to be completed, and 2438 participants met this criterion. The multilevel regression included data for which all relevant variables were available, resulting in a final sample size of 1853 patients
Contamination	Low risk	Allocation was by practice, and it is unlikely that the control practices received the intervention
Selective reporting (reporting bias)	Unclear risk	All relevant outcomes in the method section (p B) were reported in the results section (p D-E). A study protocol was not available and there was insufficient information to permit judgement of high or low risk of bias
Other bias	Low risk	Cluster-randomised trial with appropriate statistical analysis

Calland 2011

Methods	A RT of an IPC intervention aimed to determine the effectiveness of procedural checklists for surgical teams during 47 laparoscopic cholecystectomies. General surgeons were randomly assigned to an intervention (i.e. the use of the checklist) or a control group
Participants	Country: USA Ten general surgeon teams consisting of surgeons, anaesthetists and nurses. Twenty-three patients in the control group and 24 in the intervention group. Eighteen patients dropped out between the randomisation and the analysis
Interventions	An intraoperative procedural checklist including preoperative, intraoperative, and post-operative items
Outcomes	Clinical process or efficiency outcomes: length of operation, discharge status, readmission rates and technical proficiency. Collaborative behavioural outcomes: team behaviours (e.g. team communication and co-ordination)
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomisation was mentioned: "a total of 65 cases were randomized (by attending surgeon) to...", but method not specified
Allocation concealment (selection bias)	Unclear risk	No description of allocation method.
Baseline outcome measurements similar - All outcomes	Unclear risk	Not reported.
Baseline characteristics similar	Low risk	Quote: "Length of operation, discharge status, and readmission rates as indication of case outcome showed nonstatistical differences between groups."
Blinding of outcome assessment (detection bias) - All outcomes	High risk	It did not appear that there was any blinding.
Incomplete outcome data (attrition bias) - All outcomes	High risk	Acknowledged sites dropped out but ITT was not mentioned in the text Patient level: Quote: "A total of 65 cases were randomized..." Quote: "Eighteen subjects/cases dropped out between randomization and analysis: two in the checklist group declined to use the checklist or requested that their cases be withdrawn after videotaping, three cases

Calland 2011 (Continued)

		were excluded due to the conversion from laparoscopic to open procedure, procedure cancellations occurred in four cases, and scheduling difficulties or mechanical problems precluded participation for the nine remaining dropouts.”
Contamination	Unclear risk	Randomised at the level of surgeon, but as noted by the authors “there exists the possibility that residents and other staff participated in both control and intervention cases and this contaminated our results” (p 1137)
Selective reporting (reporting bias)	Unclear risk	All relevant outcomes in the method section (p 1132-3) were reported in the results section (p 1133-6). A study protocol was not available and there was insufficient information to permit judgement of high or low risk of bias
Other bias	Low risk	None detected.

Cheater 2005

Methods	A RT where 22 multidisciplinary teams from five acute care hospitals were randomised to an intervention group that participated in a facilitated programme on multidisciplinary audit or a control group	
Participants	Country: UK Nurses, physicians and other professionals (e.g. pharmacist, social worker, physiotherapist), service support staff (e.g. ward clerk, care assistant), and managers. A range of specialties (e.g. surgery, medicine, and nephrology) were included. There were 11 teams with a total of 77 participants in the intervention group and 11 teams with a total of 64 participants in the control group	
Interventions	Five facilitated meetings over 6 months with activities designed to support multidisciplinary teams to undertake an audit	
Outcomes	Collaborative audit activity.	
Notes		
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement

Random sequence generation (selection bias)	Low risk	Quote: "Teams within the same hospital were stratified on mean self-reported KSA scores, perceived level of team collaboration and medical or surgical specialty before randomisation. The project secretary under the supervision of [a researcher] randomised 22 teams to intervention or control groups, using a computer random number generator"
Allocation concealment (selection bias)	Low risk	Quote: "With the exception of two accident and emergency teams in different hospitals, teams from the same organisation were randomised in pairs. Other researchers were blind to allocation."
Baseline outcome measurements similar - All outcomes	Low risk	At baseline, both groups were equivalent for baseline variables in relation to KSA scores, and on the scores for the Collaborative Practice Scale
Baseline characteristics similar	Low risk	Quote: "At baseline, both groups were equivalent for all outcome variables except two. In comparison to the intervention group, the control arm reported higher levels of audit knowledge (median score 32.5 vs 25.0, $z = -3.001$, $P = 0.003$) and skills (median score 32.5 vs. 24.6, $z = -2.990$, $P = 0.003$). Baseline differences were adjusted for in the analysis. Baseline differences were not found for WWTs."
Blinding of outcome assessment (detection bias) - All outcomes	Low risk	Quote: "Two members of the research team (RB and HH) independently assessed the quality of the reports (blind to group allocation) and the percentage inter-rater agreement did not fall below 82%."
Incomplete outcome data (attrition bias) - All outcomes	High risk	Practice level: Quote: "Participation in the intervention programme was associated with increased audit activity, with 9 of the 11 teams reporting improvements to care and seven teams completing the full audit cycle. In contrast, the majority of teams in the control group had made no progress with undertaking an audit and only two teams had undertaken a first data collection and implemented changes."

Cheater 2005 (Continued)

		Patient level: Results were provided about the quality of the audits in relation to their compliance with the 55 quality criteria, but no further information was provided in relation to any patient level outcomes
Contamination	Low risk	Only intervention teams participated in the facilitation programme
Selective reporting (reporting bias)	Unclear risk	All relevant outcomes in the method section (p 781-2) were reported in the results section (p 785-7). A study protocol was not available and there was insufficient information to permit judgement of high or low risk of bias
Other bias	Low risk	None detected.

Curley 1998

Methods	Randomised trial - Firm trial: patients and staff from inpatient medical wards at an acute care hospital were randomised to one of six medical wards. Three wards were allocated to the intervention group that implemented daily interdisciplinary work rounds, and three wards were allocated to the control group that continued traditional work rounds
Participants	Country: USA Interns and residents in medicine, staff nurses, nursing supervisors, respirologists, pharmacists, nutritionists, and social workers. There were 567 patients in the intervention group and 535 patients in the control group
Interventions	Daily interdisciplinary work rounds.
Outcomes	Length of stay, total charges, orders for administration of aerosols
Notes	Unit of analysis error - allocated intervention to wards but analysed patients without correction for clustering. However, this correction may not substantially change the conclusion because randomisation of staff and patients limits variation between clusters

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The firm system randomization procedures and their validation have been reviewed extensively in the literature. Each inpatient firm has two physician teams or ward services. For this trial the six ward ser-

Curley 1998 (Continued)

		vices were divided so that three ward services continued traditional work rounds as usual and the three ward services implemented the CQI designed interdisciplinary work rounds.”
Allocation concealment (selection bias)	Low risk	Quote: “The firm system randomization procedures and their validation have been reviewed extensively in the literature. Each inpatient firm has two physician teams or ward services. For this trial the six ward services were divided so that three ward services continued traditional work rounds as usual and the three ward services implemented the CQI designed interdisciplinary work rounds.”
Baseline outcome measurements similar - All outcomes	Unclear risk	Not reported.
Baseline characteristics similar	Low risk	Quote: “After controlling for baseline differences in case-mix using a multivariate propensity score, the length of stay and total charges for the hospital stay for the patients included in the trial were evaluated.”
Blinding of outcome assessment (detection bias) - All outcomes	Low risk	Quote: “Patient data were retrieved from the hospital’s administrative and billing system. Thus, patient specific cost and efficiency outcomes were limited to resource utilization in the form of hospital length of stay and total charges.” “...the Respiratory Therapy (RT) Department conducted a study of aerosol use appropriateness, as determined by criteria previously devised and tested by the RT Department.”
Incomplete outcome data (attrition bias) - All outcomes	Low risk	Practice level: Quote: “The outcome measures reported in this review were at the patient level. The study does report results from satisfaction surveys completed by 19 providers of the traditional rounds group and 21 providers of the interdisciplinary rounds group but provides no information about the total number of providers in each group.” Patient level:

Curley 1998 (Continued)

		Quote: "Study patients included all patients admitted to the medical inpatient units between November 8, 1993, and May 31, 1994, who spent at least 50% of their hospital stay on that unit and were discharged from that unit. If patients were readmitted during the trial, each admission was considered separately." "Patient data were retrieved from the hospital's administrative and billing system."
Contamination	Low risk	Quote: "Patients were excluded from analysis if their hospital stay was not on their assigned medical firm because they had been 'de-firmed' because of excess admissions to one service or if they were 'boarding' on a floor that was not the ward team's home floor. Patients were excluded from the trial if they were transferred from medicine to another service (e.g. surgery) or if less than 50% of their stay occurred on the medical floor..."
Selective reporting (reporting bias)	Unclear risk	All relevant outcomes in the method section (AS6) were reported in the results section (AS7-9). There was no published protocol so we cannot be sure all planned analyses were conducted
Other bias	Low risk	None detected.

Deneckere 2013

Methods	A post-test-only cluster-RT of 30 teams caring for patients with COPD and PFF. 17 intervention teams and 13 control teams examined how the use of CPs improved teamwork in an acute hospital setting
Participants	Country: Belgium Doctors (i.e. orthopaedic surgeons or pneumologists), head nurses, nurses, and allied health professionals (i.e. physiotherapists and social workers). 581 participants: 346 in the intervention teams (N = 17) and 235 in the control teams (N = 13)
Interventions	The intervention involved the development and implementation of CPs including 3 components: 1) feedback on team's performance before CP implementation; 2) receipt of evidence-based key-indicators for implementing CPs in practice to review; 3) training in CP development. Control teams: usual care

Outcomes	Conflict management, team climate for innovation, level of organised care, emotional exhaustion, level of competence, relational co-ordination	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Stratified randomisation was used to assign the teams to an intervention group (using care pathways) and a control group (usual care). Interprofessional teams were randomised. COPD/PFF was used as blocking factor."
Allocation concealment (selection bias)	Low risk	Quote: "Before the start of the randomisation process, random numbers were assigned to each cluster by a researcher not involved in the study, using the online available tool 'Research Randomizer' www.randomizer.org). Next, the researcher randomly allocated the coded clusters to the intervention or control group using the same online tool."
Baseline outcome measurements similar - All outcomes	Unclear risk	Not reported.
Baseline characteristics similar	Low risk	Intervention and control teams were reasonably similar. Quote: "No significant differences in organizational or team member characteristics were found, except for the number of years of experience, which was significantly higher in the control group" (Table 2)
Blinding of outcome assessment (detection bias) - All outcomes	High risk	It did not appear that there was any blinding.
Incomplete outcome data (attrition bias) - All outcomes	Low risk	Practice level ITT was not mentioned. Authors acknowledged that sites dropped out Quote: "A potential weakness of the study is the dropout of 7 teams and its possible impact on the results."
Contamination	Low risk	Only intervention teams participated in the development and implementation of CP

Deneckere 2013 (Continued)

Selective reporting (reporting bias)	Low risk	All relevant outcomes in the method section (p 100-1) were reported in the results section (p 102-4). There was also a published protocol and all planned analyses were conducted
Other bias	Low risk	Cluster-RT with appropriate statistical analysis.

Schmidt 1998

Methods	A RT of 33 nursing homes, 15 experimental homes and 18 control homes, to examine the effects of monthly facilitated multidisciplinary rounds on the quality and quantity of psychotropic drug prescribing.	
Participants	Country: Sweden Physician, pharmacists, selected nurses, and nursing assistants 1854 long-term residents: 626 in experimental homes and 1228 in control homes	
Interventions	Pharmacist led team meetings once a month over a period of 12 months	
Outcomes	Proportion of patients receiving drugs, number of psychotropic drugs, use of non-recommended hypnotics, use of non-recommended anxiolytics, use of non-recommended antidepressant drugs	
Notes		

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not reported.
Allocation concealment (selection bias)	Low risk	Quote: "Thirty-six nursing homes, representing 5% of all nursing homes in Sweden, participated in the study. The sampling process consisted of three steps. At the time of the study, the National Corporation of Swedish Pharmacies was organized into 36 regions, 18 of which were randomly selected for this study. Each regional pharmacy director then selected two facilities in his or her region using several criteria....Researchers randomly assigned one home in each pair to receive the intervention."

Schmidt 1998 (Continued)

Baseline outcome measurements similar - All outcomes	Low risk	Quote: "At baseline, we found no significant differences in the proportion of residents with scheduled psychotropics (64% vs 65%), number of drugs among residents with psychotropics (2.07 vs 2.06)."
Baseline characteristics similar	Low risk	Quote: "There were no significant differences in the demographic, functional, or psychiatric characteristics of residents in experimental and control homes at baseline." Quote: "The overall level of prescribing was similar in experimental and control homes before the intervention (Table 2). At baseline, we found no significant differences in the proportion of residents with scheduled psychotropics (64% vs 65%), number of drugs among residents with psychotropics (2.07 vs 2.06), or proportion of residents with polymedicine (46% vs 47%). Baseline rates of therapeutic duplication were also comparable in the experimental and control homes."
Blinding of outcome assessment (detection bias) - All outcomes	Low risk	Quote: "Lists of each resident's prescriptions were collected 1 month before and 1 month after the 12-month intervention in both experimental homes and control homes. Trained coders, supervised by pharmacists, classified and coded all scheduled and PRN (pro re nata) orders."
Incomplete outcome data (attrition bias) - All outcomes	Low risk	3 intervention homes out of 18 became ineligible.
Contamination	Low risk	Quote: "Pharmacists assigned to experimental homes had no contact with control nursing homes. In the control homes, no efforts were made beyond normal routine to influence drug prescribing."
Selective reporting (reporting bias)	Unclear risk	Insufficient information to permit judgement of high or low risk of bias. There was no published protocol so we cannot be sure all planned analyses were conducted
Other bias	Low risk	None detected.

Strasser 2008

Methods	RT, in which patients with a stroke were treated by 31 teams from 31 Veteran Affairs rehabilitation units before and after a multifaceted intervention, aimed at improving interprofessional collaboration
Participants	Country: USA Medical doctors, nurses, occupational therapists, speech-language pathologists, physical therapists, and case managers or social workers. 464 participants: 227 in the intervention teams (N = 15) and 237 in the control teams (N = 16). Patients with a stroke were randomly assigned to each group
Interventions	Intervention teams: received the following multifaceted intervention: 1) an off-site workshop emphasising team dynamics, problem-solving, and the use of performance feedback data; 2) action plans (specific team performance profiles with recommendations) for process improvement; 3) telephone and video conference consultations to sustain improvement in collaboration Control teams only received specific team performance profile Information
Outcomes	Functional improvement (as measured by the change in motor items of the FIM instrument), length of stay (LOS), rates of community discharge

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "... we randomized sites to either intervention or control group using a computer; each stratum was force randomized to have 4 sites in 1 arm."
Allocation concealment (selection bias)	Unclear risk	Method of concealment was not described.
Baseline outcome measurements similar - All outcomes	Low risk	The mean FIM scores at baseline were similar for the intervention group (52.2 ± 3.9) and for the control group (52.4 ± 3.8)
Baseline characteristics similar	Low risk	Quote: "...There were no differences between study conditions in demographic characteristics (table 2). Control sites admitted stroke patients with lower initial (admission) motor FIM scores during the pre-intervention periods (P.002); thus, we adjusted all analyses using FRGs ... a classification based on initial motor FIM and age."
Blinding of outcome assessment (detection bias) - All outcomes	High risk	It did not appear that there was any blinding.

Strasser 2008 (Continued)

Incomplete outcome data (attrition bias) - All outcomes	High risk	Acknowledged sites dropped out but ITT was not mentioned in the text Practice level Quote: "Of 33 eligible sites, a total of 31 sites agreed to participate, initiated the IRB approval, and were randomized. One control site was unable to complete the IRB process and withdrew, and 1 intervention site did not report data to the FSOD, leaving 15 sites in the control group and 14 in the intervention group"
Contamination	Low risk	No reason to think contamination had occurred.
Selective reporting (reporting bias)	Low risk	All relevant outcomes in the methods section (p 11) were reported in the results section (p 14). There was a published protocol and all planned analyses were conducted
Other bias	Low risk	None detected.

Wild 2004

Methods	Randomised trial in which patients in inpatient telemetry ward in a community hospital were randomised to the intervention medical team, which conducted interdisciplinary rounds or to the control team, which provided standard care	
Participants	Country: USA Resident physicians, nurses, a case manager, pharmacist, dietician, and physical therapist. Eighty-four patients were enrolled: 42 in intervention and 42 in standard care	
Interventions	Intervention: daily interdisciplinary rounds. Control group: standard care.	
Outcomes	Length of hospital stay	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Randomization was performed using random numerical assignments in pre-sealed envelopes."

Allocation concealment (selection bias)	Unclear risk	Envelope randomisation.
Baseline outcome measurements similar - All outcomes	Low risk	Mean length of stay (days) was similar in the intervention group (3.04 ± 1.8) compared with the control group (2.7 ± 1.8)
Baseline characteristics similar	Low risk	Quote: "There were no significant differences between groups for admission diagnosis; number of co-morbidities; number of abnormal laboratory data; ability to perform activities of daily living; presence of dementia or diabetes, or whether there was a home health aide. In spite of randomization, the gender composition between groups was somewhat different...and the number of readmissions in the IR Team was higher than in the non-IR Team ($P = 0.003$)."
Blinding of outcome assessment (detection bias) - All outcomes	Low risk	Quote: "Charts were surveyed to determine patient characteristics and LOS. LOS was measured as the difference between discharge and admission date."
Incomplete outcome data (attrition bias) - All outcomes	Low risk	Practice level: Quote: "Questionnaire return was 80%", but these results were not reported in this review because they did not meet outcome criteria Patient level: All participants were accounted for and none were lost to follow-up
Contamination	Low risk	Quote: "Patients were randomly assigned to two medical teams: the intervention group received IRs and the control subjects received standard care."
Selective reporting (reporting bias)	Unclear risk	All relevant outcomes in the method section (p 64) were reported in the results section (p 67). There was no published protocol so we cannot be sure all planned analyses were conducted
Other bias	Low risk	None detected.

Wilson 2004

Methods	RT comparing multidisciplinary audio conferencing and multidisciplinary video conferencing with a team that worked at two hospitals
Participants	Country: Australia Medical staff specialists, medical registrars, nurses, speech pathologist, occupational therapists, social worker, medical students. Fifty patients were randomly assigned to each group
Interventions	Multidisciplinary audio conferences and video conferences. At each conference session, the audio conferences were conducted before the video conferences, with the same multidisciplinary team
Outcomes	Number of audio conferences held per patient, number of video conferences held, length of treatment

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The random allocation was done by an independent administrative assistant, using a table of random numbers."
Allocation concealment (selection bias)	Low risk	Quote: "The random allocation was done by an independent administrative assistant, using a table of random numbers."
Baseline outcome measurements similar - All outcomes	Unclear risk	None reported.
Baseline characteristics similar	Low risk	Quote: "The two groups were similar in terms of age, sex and diagnosis (Table 1)."
Blinding of outcome assessment (detection bias) - All outcomes	Low risk	Quote: "Conference times were recorded by an independent observer and files were reviewed by an independent medical practitioner blinded to the randomization."
Incomplete outcome data (attrition bias) - All outcomes	Unclear risk	Practice level: Quote: "Only 14 of 29 (including 6 medical students) completed a staff satisfaction survey. These results are not reported in this review because they did not meet outcome criteria." Patient level" Quote: "There were no deaths, and all patients recruited completed the trial."

Contamination	Unclear risk	Quote: “Within each meeting of the multidisciplinary team, the audioconferences were conducted before the videoconferences, to ensure that there was no visual contact between the two locations until the latter part of the session.” “The team remained consistent at either site for both the audio- and videoconferences held on each individual day of the conference, but the team members rotated between sites over the study period.” While measures were taken to prevent contamination, the same team members were involved in both types of conferencing
Selective reporting (reporting bias)	Unclear risk	All relevant outcomes in the method section (p 353-4) were reported in the results section (p 354). Insufficient information was provided to permit judgement of high or low risk of bias. There was no published protocol so we cannot be sure all planned analyses were conducted
Other bias	Low risk	None detected.

CCTP = Chronic care team profile
 COPD = chronic obstructive pulmonary disease
 CP = care pathway
 CQI = Continuous quality improvement
 FIM = Functional independence measure
 FRG = Functional-related groups
 FSOD = Functional status outcomes database
 GP = General practitioner
 IPC = Interprofessional collaboration
 IRs = interdisciplinary rounds
 IRB = Institutional Research Board
 ITT = Intention-to-treat
 KSA = Knowledge, skills, attitudes
 LOS = length of stay
 PACIC = Patient assessment of chronic illness care
 PFF = proximal femur fracture
 RT = randomised trial
 WWT = Wider ward teams

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Bekelman 2015	Not a practice-based IPC intervention
Boet 2013	Not a practice-based IPC intervention
Boone 2008	Not a practice-based IPC intervention
Chen 2013	Not a practice-based IPC intervention
Cheng 2013	Not a practice-based IPC intervention
Curtis 2012	Not a RT
Dhalla 2014	Not a practice-based IPC intervention
Döpp 2013	Not a practice-based IPC intervention
Fransen 2012	Not a practice-based IPC intervention
Goud 2009	Not a practice-based IPC intervention
Hallin 2011	Not a practice-based IPC intervention
Hobgood 2010	Not a practice-based IPC intervention
Hoffmann 2014	Not a practice-based IPC intervention
Jankouskas 2011	Not a practice-based IPC intervention
Jenkins 2013	Not a practice-based IPC intervention
Katakam 2012	Not a practice-based IPC intervention
Keller 2012	Not a practice-based IPC intervention
Kemper 2011	Not a RT
Koerner 2014	Not a practice-based IPC intervention
Kunkler 2007	Not a practice-based IPC intervention
Körner 2012	Not a practice-based IPC intervention
Lee 2013	Not a practice-based IPC intervention

(Continued)

Marsteller 2013	Not a practice-based IPC intervention
Mohaupt 2012	Not a practice-based IPC intervention
Musick 2011	Not a practice-based IPC intervention
O'Leary 2010	Not a RT
Rörtgen 2013	Not a practice-based IPC intervention
Van de Ven 2010	Not a RT
Weller 2014	Not a practice-based IPC intervention
Wittenberg-Lyles 2013	Not a practice-based IPC intervention

IPC: interprofessional collaboration

RT: randomised trial

DATA AND ANALYSES

This review has no analyses.

APPENDICES

Appendix 1. CENTRAL search strategy

```
#1[mh "Interprofessional Relations"] and (collaborat* or team*)
#2 [mh "Patient Care Team"] and (collaborat* or team*)
#3 ((interprofession* or inter-profession*) next (collaborat* or team*))
#4 ((interdisciplin* or inter-disciplin*) next (collaborat* or team*))
#5 ((interoccupation* or inter-occupation*) next (collaborat* or team*))
#6 ((multiprofession* or multi-profession*) next (collaborat* or team*))
#7 ((multidisciplin* or multi-disciplin*) next (collaborat* or team*))
#8((multioccupation* or multi-occupation*) next (collaborat* or team*))
#9((transdisciplin* or trans-disciplin*) next (collaborat* or team*))
#10(team* next collaborat*)
#11{or #1-#10}
```

Appendix 2. MEDLINE search strategy

```
1 exp Interprofessional Relations/ and (collaborat$ or team$).tw. (8220)
2 exp Patient Care Team/ and (collaborat$ or team$).tw. (13439)
3 ((interprofession$ or inter-profession$) adj (collaborat$ or team$)).tw. (853)
4 ((interdisciplin$ or inter-disciplin$) adj (collaborat$ or team$)).tw. (2660)
5 ((interoccupation$ or inter-occupation$) adj (collaborat$ or team$)).tw. (0)
6 ((multiprofession$ or multi-profession$) adj (collaborat$ or team$)).tw. (355)
7 ((multidisciplin$ or multi-disciplin$) adj (collaborat$ or team$)).tw. (7856)
8 ((multioccupation$ or multi-occupation$) adj (collaborat$ or team$)).tw. (0)
9 ((transdisciplin$ or trans-disciplin$) adj (collaborat$ or team$)).tw. (105)
10 (team$ adj collaborat$).tw. (158)
11 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 (26183)
12 randomized controlled trial.pt. (276233)
13 controlled clinical trial.pt. (42446)
14 randomized controlled trials/ (83918)
15 random allocation/ (45887)
16 double blind method/ (80591)
17 single blind method/ (16519)
18 12 or 13 or 14 or 15 or 16 or 17 (452549)
19 animals/ not humans/ (1793680)
20 18 not 19 (410246)
21 11 and 20 (954)
22 limit 21 to yr="2007 -Current" (595)
```

Appendix 3. CINAHL search strategy

1 (MH "Interprofessional Relations+") AND TX ((collaborat* or team*))
2 (MH " Multidisciplinary Care Team+") AND TX ((collaborat* or team*))
3 TX ((interprofession* or inter-profession*) N1 (collaborat* or team*))
4 TX ((interdisciplin* or inter-disciplin*) N1 (collaborat* or team*))
5 TX ((interoccupation* or inter-occupation*) N1 (collaborat* or team*))
6 TX ((multiprofession* or multi-profession*) N1 (collaborat* or team*))
7 TX ((multidisciplin* or multi-disciplin*) N1 (collaborat* or team*))
8 TX ((multioccupation* or multi-occupation*) N1 (collaborat* or team*))
9 TX ((multioccupation* or multi-occupation*) N1 (collaborat* or team*))
10 TX ((transdisciplin* or trans-disciplin*) N1 (collaborat* or team*))
11 TX team* N1 collaborat*
12 S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11
13 (MH "Clinical Trials+")
14 PT Clinical trial
15 TX ((trebl* n1 blind*) or (trebl* n1 mask*))
16 TX ((tripl* n1 blind*) or (tripl* n1 mask*))
17 TX ((doubl* n1 blind*) or (doubl* n1 mask*))
18 TX ((singl* n1 blind*) or (singl* n1 mask*))
19 TX randomi* control* trial*
20 (MH "Random Assignment")
21 TX random* allocat*
22 TX placebo*
23 (MH "Placebos")
24 (MH "Quantitative Studies")
25 TX allocat* random*
26 S13 OR S14 OR S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24 OR S25
27 S12 AND S26
28 S12 AND S26. Limiters - Publication Year: 2007-2014; Clinical Trial

Appendix 4. ClinicalTrials.gov and ICTRP search strategies

ClinicalTrials.gov search strategy

(collaboration OR team) AND (interdisciplinary OR interprofessional OR multidisciplinary OR multiprofessional)

ICTRP search strategy

#1 collaboration AND interdisciplinary
#2 collaboration AND interprofessional
#3 collaboration AND multidisciplinary
#4 collaboration AND multiprofessional
#5 team AND interdisciplinary
#6 team AND interprofessional
#7 multidisciplinary team
#8 team AND multiprofessional
#9 OR/1-8

WHAT'S NEW

Last assessed as up-to-date: 10 November 2015.

Date	Event	Description
13 November 2015	New citation required but conclusions have not changed	This update found four new studies. As a result, the review now includes nine studies. While the number of studies has increased slightly, the main conclusions from the previous update remain unchanged (Zwarenstein 2009). There have been changes to the author team, with the inclusion of two new authors
10 November 2015	New search has been performed	New searches performed to 10 November 2015. Four new studies identified

HISTORY

Protocol first published: Issue 3, 1996

Review first published: Issue 2, 1997

Date	Event	Description
13 May 2009	New citation required and conclusions have changed	Conclusions changed, based on additional studies. Criteria for included study designs, included participants and specification of the intervention changed from the 1997 review. This first review included randomised trials, controlled before-after studies and interrupted time series designs, whereas this update included only randomised trials. The types of participants included in the first review were physicians and nurses, whereas this update included all types of healthcare professionals. The first review included studies in which the interventions may not have specified their intent to change interprofessional collaboration, whereas this update only included studies with an explicit focus on collaboration. These changes were intended to increase the validity of the conclusions, and to widen their applicability to professions other than nursing and medicine
13 May 2009	New search has been performed	New search and four additional studies identified and included in the review
20 August 2008	New search has been performed	Converted to new review format.
11 January 2000	New citation required and conclusions have changed	Substantive amendment

CONTRIBUTIONS OF AUTHORS

SR and FP undertook the searches and both independently reviewed each of the titles and abstracts located to ensure they met the inclusion criteria. SR and FP independently assessed each full-text article to ensure they met the criteria. MZ resolved any disagreements during the screening processes. SR and MZ independently assessed the certainty of evidence for the included studies. All authors (SR, FP, RH, JG, MZ) analysed and interpreted the data and contributed to writing the review. SR is guarantor for the review.

DECLARATIONS OF INTEREST

Scott Reeves: none known.

Ferruccio Pelone: none known

Reema Harrison: none known

Joanne Goldman: none known

Merrick Zwarenstein: none known.

The authors have no personal or professional interests as to whether this review shows benefits of practice-based interventions on interprofessional collaboration.

SOURCES OF SUPPORT

Internal sources

- Faculty of Health, Social Care and Education, Kingston University and St George's, University of London, UK.
- Continuing Education and Professional Development, Faculty of Medicine, University of Toronto, Canada.

External sources

- Canadian Institutes of Health Research, Canada.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Review authorship changed between protocol and review updates. The title has changed from the protocol: 'The effects on patient care of interventions to change collaboration between nurses and doctors' (Zwarenstein 1996). Changes were also made to criteria for included study designs, included participants, and specification of the intervention between the published protocol and this update. The protocol planned to include randomised trials, controlled before-after studies, and interrupted time series designs, whereas the last update (Zwarenstein 2009) and this update included only randomised trials. The protocol planned to include only physicians and nurses, whereas this update included all types of health and social care professionals. The protocol also planned to include studies in which the interventions may not have specified their intent to change interprofessional collaboration, whereas this update included only studies with an explicit focus on collaboration. These changes were made to increase the validity of the conclusions, and to widen their applicability to professions beyond nursing and medicine.

INDEX TERMS

Medical Subject Headings (MeSH)

*Cooperative Behavior; *Health Personnel; *Interprofessional Relations; *Professional Practice; Delivery of Health Care; Quality of Health Care; Randomized Controlled Trials as Topic; Telecommunications

MeSH check words

Female; Humans