

©2020. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/about/downloads>



This is the accepted version of this article. The version of record is available at:
<https://doi.org/10.1016/j.foodcont.2020.107852>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

**Blueberry, Raspberry, and Strawberry Extracts Reduce the Formation of
Carcinogenic Heterocyclic Amines in Fried Camel, Beef and Chicken meats.**

Mohammad Rizwan Khan¹, Rosa Busquets*², Mohammad Azam¹

¹Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi
Arabia

²School of Life Sciences, Pharmacy and Chemistry, Kingston University London, Penrhyn
road, KT1 2EE, Kingston upon Thames, United Kingdom

Corresponding author.

E-mail address: r.busquets@kingston.ac.uk (R. Busquets).

28 **Abstract**

29 Heterocyclic amines (HCAs) are toxic products from the Maillard reaction that form from
30 the reaction of sugars, amino acids and creatine/creatinine when cooking protein rich food.
31 In this work, commonly consumed meats in Saudi Arabia (camel, beef and chicken) were
32 fried under conditions resembling home cooking. The effect of marinades made of
33 blueberry, raspberry and strawberry were tested separately on meat at different marinating
34 times (1, 6, 12, 24h, at 4°C) before frying. The marinades caused an overall reduction of
35 HCAs. The decrease was more noticeable with long marination time ≥ 6 h. The reduction of
36 individual HCAs, after 24h marinades, was 91-100% for pyridines; 40-67% for β -
37 carbolines; and 100% for quinoxalines, quinolines, α -carbolines and γ - carbolines, although
38 the latter three were seldomly detected in this study. An increase, up to 2 times, on the
39 formation of the studied quinoxalines was observed in every meat and marination for no
40 more than 1h. Therefore, longer marinating times with berry extracts, from 6h, are
41 recommended over those below (1h).

42

43 **Keywords:** marinade; PhIP; UPLC-MS/MS; Maillard reaction; foodborne carcinogen

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 Highlights

- 60 • From high to low concentration of mutagenic HCAs: fried chicken, camel and beef
- 61 • 1h blueberry, raspberry and strawberry juice marinades boosted quinoxalines
- 62 • ≥ 6 h blueberry, raspberry and strawberry marinades reduced quinoxaline HCA levels
- 63 • ≥ 12 h marinades had high impact on the reduction of pyridines and β -carbolines
- 64 • 24h marinades caused 40-100% reduction in total HCA

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99 **1. Introduction**

100

101 In the last 30 years, the occurrence of the foodborne carcinogenic heterocyclic amines
102 (HCAs) in various protein-rich cooked foods such as meat and fish has been extensively
103 investigated (Barzegar, Kamankesh & Mohammadi, 2019, Khan, Busquets, Saurina,
104 Hernández, S., & Puignou, 2013, Lu, Kuhnle, & Cheng, 2017). Thus far, over 24 HCAs
105 have been identified in cooked food and it is accepted that HCAs can form from reactions of
106 amino acids, creatine/creatinine and sugar, although these 3 types of biomolecules are not
107 essential for the formation of all HCAs (Skog, Johansson & Jägerstad 1998; Murkovic, 1999
108 ; Gibis & Weiss, 2015). Structurally, HCAs found in food are in the form of
109 aminocarboline and aminoimidazoazaarenes. While aminocarboline are described to form
110 from amino acids and protein pyrolysis at high temperatures (>300°C),
111 aminoimidazoazaarenes form readily at lower temperatures via aldol condensation of
112 pyrazines or pyridines with aldehydes and creatinine (Naushad & Khan, 2014; Oz & Kotan,
113 2016).

114 The relationship between the consumption of red meat and the likelihood of developing
115 different types of cancer has been established in epidemiological studies (Oostindjer et al.,
116 2014), however the link between exposure to HCAs and the onset of these cancers remains
117 unclear (Bellamri & Turesky, 2019). Animals studies and clinical trials have been
118 performed to elucidate the causative link between exposure to HCAs and alterations in DNA
119 (Turesky & Vouros, 2004; Tang, Kassie, Qian, Ansha & Turesky, 2013). However, many of
120 the existing studies were carried out with HCAs concentrations and exposure-times that do
121 not resemble those in a normal diet (Felton et al., 2007).

122 Recent studies have revealed a correlation between the intake of the HCA 2-amino-1-
123 methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) and the likelihood of developing cancer
124 (Rogers et al., 2016, Bellamri, Xiao, Murugan, Weight & Turesky, 2018). Indeed, several
125 HCAs are categorized as possible or probable human carcinogens by the International
126 Agency for Research on Cancer (IARC), and there is a recommendation for a reduction of
127 their consumption (IARC, 1993). The US National Toxicology Program (NTP) also listed
128 some HCAs as reasonably anticipated human carcinogens (NTP, 2004). The discovery of
129 mutagenic forms of HCAs and their adducts with DNA in human tissues is indicative of
130 their toxicity under common meat intake levels through diet (Busquets, Frandsen, Jönsson,
131 Puignou & Galceran & Skog 2013, Bellamri, Xiao, Murugan, Weight & Turesky, 2018,
132 Guo et al., 2018).

133 However, the exposure to HCA is not unavoidable. The intake of HCAs' through the
134 consumption of meat and fish can be reduced by adopting particular cooking practices such as
135 reducing cooking temperature and time, decreasing superficial cooking temperature with
136 water (e.g. stews) and using ingredients that affect the transport of HCAs' precursors to the
137 food surface, where temperature will be greater. In this regard, the addition of ingredients
138 with water-holding capacity or marinating methods have been shown to be effective at
139 reducing the formation of HCAs (Persson, Sjöholm & Skog, 2003; Vitaglione & Fogliano,
140 2004; Oz & Kaya, 2011).

141 In Saudi Arabia, HCAs have been reported in camel (Khan, Naushad & Zeid, 2017) and
142 chicken items from local restaurants (Alsohaimi, Khan, Ali, & Azam, 2019), with some
143 chicken dishes presenting relatively high levels of MeIQx (2-3 ng/g) and PhIP (7-36 ng/g)
144 compared to their levels reported in other items (Busquets, 2012). Recipes including
145 marinades could have an important impact on the formation of HCAs due to the presence of

146 radical scavengers but also the effect of sugars, pH and the aqueous environment that will
147 affect the transport of HCA's precursors within meat. The main hypothesis of this study is
148 that fruit-based marinades (blueberry, raspberry and strawberry) can be effective at reducing
149 the formation of HCAs during the cooking of camel, beef and chicken, three types of meat
150 that are highly consumed in Saudi Arabia but also elsewhere.

151

152 **2.1. Materials and chemicals**

153 Acetonitrile, ethyl acetate and methanol of LC grade were obtained from Merck
154 (Darmstadt, Germany). Ammonium acetate ($\geq 98\%$), ammonium formate ($\geq 99\%$), ammonia
155 solution (25%) formic acid ($\geq 98\%$) and NaOH ($\geq 97\%$) were purchased from Merck
156 (Darmstadt, Germany). Fifteen HCAs (structures given in Figure 1S) were studied: 2-
157 amino-1,6-dimethylimidazo[4,5-*b*]pyridine (DMIP), 2-amino-1-methyl-6-
158 phenylimidazo[4,5-*b*]pyridine (PhIP), 2-amino-3-methylimidazo[4,5-*f*]quinoline (IQ), 2-
159 amino-3,4-dimethylimidazo[4,5-*f*]quinoline (MeIQ), 2-amino-3,8-dimethylimidazo[4,5-
160 *f*]quinoxaline (MeIQx), 2-amino-3,4,8-trimethylimidazo[4,5-*f*]quinoxaline (4,8-DiMeIQx),
161 2-amino-3,7,8-trimethylimidazo[4,5-*f*]quinoxaline (7,8-DiMeIQx) 2-amino-3,4,7,8-
162 tetramethylimidazo[4,5-*f*]quinoxaline (4,7,8-TriMeIQx, internal standard), 2-amino-6-
163 methyl-dipyrido [1,2-*a*:3',2'-*d*]imidazole (Glu-P-1), 2-amino- dipyrido[1,2-*a*:3'2'-
164 *d*]imidazole (Glu-P-2), 2-amino-9*H*-pyrido[2,3-*b*]indole (A α C), 2-amino-3-methyl-9*H*-
165 pyrido[2,3-*b*]indole (MeA α C), 3-amino-1,4-dimethyl-5*H*-pyrido[4,3-*b*]indole (Trp-P-1), 3-
166 amino-1-methyl-5*H*-pyrido[4,3-*b*]indole (Trp-P-2). These HCAs were obtained from
167 Toronto Research Chemicals (Toronto, Canada). The co-mutagenic amines 1-methyl-9*H*-
168 pyrido[3,4-*b*]indole (harman) and 9*H*-pyrido[3,4-*b*]indole (norharman) were purchased

169 from Sigma-Aldrich (Missouri, USA). The HCAs purity was >99%. 4,7,8-TriMeIQx was
170 added in standards and purified sample extracts as internal standard.

171 The HCAs stock standard solutions were prepared at 200 µg/mL in methanol and used for
172 spiking samples in standard addition. Calibration curves with standard mixtures of fifteen
173 HCAs between 0.001 µg HCAs/mL and 1.00 µg HCAs/mL were prepared to establish the
174 linearity range. 4,7,8-TriMeIQx was added in every standard at constant concentration.
175 Both standard and sample extracts were filtered using a 0.22 µm polytetrafluoroethylene
176 (PTFE) syringe filters (Macherey-Nagel, Düren, Germany) before being injected into the
177 ultra-performance liquid chromatography (UPLC) system.

178

179 2.2. Meat sample preparation and cooking

180 Fresh meat (camel loin, beef fillet and chicken breast) and cooking ingredients (blueberry,
181 raspberry, strawberry and olive oil) were purchased in a local store (Riyadh, Saudi Arabia).

182 The meat and oil were locally produced and berries, which trademark was Driscoll's, were
183 imported: from Mexico (blueberries) and the US (raspberries and strawberries). The visible
184 fat in the meat, including chicken skin, was removed and the meat was cut into fillets of
185 nearly 1 cm in thickness. Blueberries, raspberries and strawberries, individually, were
186 washed with water, cut into small pieces; blended with a juice extractor (Kenwood JE730,
187 China) and filtered to remove pulps and fibres. Individual meat fillets (100 g) and fruit
188 extracts (100 mL) were marinated at different time periods (1, 6, 12 and 24h), at 4 °C, to
189 avoid any microbial contamination. A set of unmarinated samples were used as control
190 samples. Both the marinated and unmarinated meat samples were pan-fried.

191 A gas cooker (Gibson, Cairo, Egypt) and a non-stick frying pan (Tefal, Durbase
192 Technology, Paris, France) was used. The cooking temperature of the meat samples was

193 measured with type K probes and TC6 software (Nomadics Inc., Stillwater, Oklahoma,
194 USA). Prior to the cooking of meat samples, the probes were calibrated by submerging
195 them in boiling water (Milli-Q) and readings adjusted to 100 °C. Cooking temperature was
196 monitored and recorded every five seconds. The European Prospective Investigation into
197 Cancer and Nutrition (EPIC) defines frying cooking method as cooking of food in either fat
198 or oil. In this study, to prevent the meat sticking to the pan, 5 mL of olive oil was added to
199 the pan at the beginning of the cooking process. The cooking started when the temperature
200 in the centre of the pan with a layer of oil was between 215°C and 230 °C. The total
201 cooking time was eight minutes: the meats were moved around the with oil for 4 minutes,
202 following which they were flipped and moved around the pan for 4 min more.
203 Subsequently, the cooked meat samples were cleaned and all pan residues, including
204 retained oil, were removed. Cooking weight loss was measured by weighing the meat
205 before and after cooking. Every meat fillet was marinated and cooked independently in
206 duplicate. Control samples were also prepared in duplicate. The meat crusts from the
207 cooked meet were separated, pooled, ground and refrigerated until characterisation. Meat
208 samples were blended using a stardust coffee grinder, CML-1000MKII (Osaka, Japan), and
209 a Microtron® MB800, Kinematica AG (Littau, Switzerland).

210

211 2.3. HCAs extraction from meat samples and quantification.

212 The cartridges used for the extraction of HCAs were octadecylsilane (C₁₈, 100 mg) and
213 Bond Elut propylsulfonyl silica (PRS, 500 mg). These solid phase extraction (SPE)
214 cartridges, connectors and stopcocks were obtained from Varian (Harbor City, USA).
215 Extraction columns (Extrelut NT20) were purchased from Merck (Darmstad, Germany).
216 Hydromatrix bulk material (diatomaceous earth) was purchased from Agilent Technologies

217 (Santa Clara, California, USA). The SPE was carried out with Visiprep™ and Visidry™
218 vacuum manifolds, from Supelco (Gland, Switzerland). They were used for the purification
219 of HCAs, and drying the elution solvent through evaporation, respectively.

220 The refrigerated ground meat crusts were allowed to equilibrate at room temperature (25
221 °C) for >30 min. Sodium hydroxide solution (50 mL, 1M) was added to the ground meat
222 crusts (20 g) followed by homogenization using ultra-turrax T25 digital homogenizer (from
223 IKA®-WERKE GmbH, Staufen, Germany). Homogenised meat samples (3 g) were
224 carefully mixed with hydromatrix bulk material (14 g, diatomaceous earth) and moved to
225 an empty column (60 mL) connected to PRS cartridge (500 mg).

226 The PRS cartridge was previously preconditioned using HCl 0.1 M (5 mL), water (10 mL)
227 and methanol (5 mL). Ethyl acetate (75 mL) was used to extract the HCAs from the
228 homogenized samples dispersed in diatomaceous earth and these were eluted to the PRS
229 cartridge. After the elution, the PRS cartridge was dried under vacuum and washed
230 sequentially using MilliQ water and methanol (4:6, v/v, 15 mL), and Milli Q water (5 mL).

231 The PRS cartridge was then coupled to a C₁₈ cartridge (100mg) which had been
232 preconditioned using methanol (5 mL) and Milli Q water (5 mL). The HCAs were eluted
233 from PRS cartridge to C₁₈ cartridge using ammonium acetate (0.5 M, pH 8.5, 20 mL). As a
234 final step, the C₁₈ cartridge was washed using Milli Q water (5 mL) followed by drying
235 under low vacuum. The HCAs elution from C₁₈ cartridge to a microcentrifuge tube was
236 performed using a methanol and ammonia solution (9:1, v/v, 800 µL). The sample solvent
237 was vaporized mildly using nitrogen. The dried sample extract was reconstituted in
238 methanol containing internal standard (4,7,8-TriMeIQx, 0.5 µg/g, 100 µL). After the
239 reconstitution, the samples were filtered (syringe filter PTFE, 0.22µm) and resolved by
240 UPLC tandem mass spectrometry (UPLC–MS/MS).

241

242 The quantification of HCAs in meat samples was carried out by standard additions method,
243 which consisted of adding a mixture of HCAs at three levels of concentration (50%, 100%
244 and 200%) with respect to the estimated initial level of HCAs in the sample. A duplicate of
245 the sample was processed and analysed without having been spiked. Specifically, The
246 samples were spiked with DMIP, PhIP, IQ, MeIQ, Glu-P-1, Glu-P-2, A α C, MeA α C, Trp-P-
247 1 and Trp-P-2 at final concentration levels of 0, 10, 50, and 150 ng HCAs/meat g and for
248 harman, norharman, MeIQx, 4,8-DiMeIQx and 7,8-DiMeIQx were 0, 5, 10 and 30 ng
249 HCAs/ g meat. The standard addition quantification of every type cooked meat was carried
250 out in triplicate. Recovery rates were estimated from the slope of the linear regression
251 between the added and recovered HCAs amounts in the meat samples.

252 2.4. Instrumentation

253 2.4.1. HCAs separation

254 The optimal HCAs separation was performed using an UPLC (Acquity®, Waters, Milford,
255 USA). The analytical column used was an ethylene bridged hybrid (BEH C₁₈) with (50 mm
256 × 2.1 mm i.d. and 1.7 μ m particle size, Acquity® from Waters (Milford, USA). The mobile
257 phase used was acetonitrile (A) and buffer solution (30 mM formic acid/ammonium
258 formate, pH 4.7, B) at 500 μ L/min. The elution programme was: 5% A in B; 0–0.1 min; 5–
259 30% A in B, 0.1–1.5 min; 30–60% A in B, 1.5–1.8 min; 60% A in B, 1.8–2.5 min. As
260 precaution, the column was washed for 2 min with methanol:water (50:50) every twenty
261 sample injections. The injection volume was 5 μ L. This analytical method was adopted
262 from a previously developed method (Barcelo-Barrachina, Moyano, Galceran, Lliberia,
263 Bago & Cortes, 2006), with minor changes.

264 2.4.2. HCAs determination

265 The HCAs were detected with a triple quadrupole mass analyser model Quattro Premier
266 Micromass (Milford, USA) equipped with electrospray (ESI) working in positive mode.
267 The quantification was carried out in multiple reaction monitoring (MRM) mode. The
268 protonated HCA molecular ions $[M+H]^+$ were the precursor ions that were fragmented to
269 product ions that were used for the quantification and confirmation of the analytes (see
270 Table 1). The working conditions of the ESI source were: 100 °C source temperature; 350
271 °C desolvation temperature; 3.6 KV capillary voltage; 38 V cone voltage; 700 L/h
272 desolvation gas; 70 L/h cone gas. High purity of nitrogen gas was used, produced from
273 Peak Scientific nitrogen generator (NM30LA, Inchinnan, United Kingdom) for the for the
274 cone gas. High purity argon for the collision gas was from Speciality Gas Centre, (Jeddah,
275 Saudi Arabia). The software used for the analysis was Waters MassLynx V4.1 (Milford,
276 USA).

277 2.4.3. Statistical analysis

278 The comparison of the concentration of HCAs with marinating time and berry extrats was
279 carried out with 2-way ANOVA with replicates and student-t test comparing means using
280 Microsoft™ Excel 2019.

281

282 **3. Results and Discussion**

283

284 Marinating meat prior cooking has shown to be among the most effective ways to reduce
285 the overall formation of HCAs (Busquets, Puignou, Galceran & Skog, 2006; Manful et al.,
286 2020). This is due to both physical and chemical effects of marinades on the Maillard
287 reaction leading to the formation of HCAs. This study explores whether HCAs levels in
288 commonly consumed meat can be reduced effectively with fruit extracts. The fruit extracts

289 tested here have potential to affect the formation of HCAs and they can be used in recipes
290 that consumers may accept.

291 The composition of the marinades was chosen on the basis of well accepted health benefits
292 of the studied berry marinades (Gowd, Bao & Chen, 2019; Zhou, Xie, Yang & Liu, 2020).

293 The cooking was carried out with full control of temperature. An example of the
294 temperature profile is given in Figure 1. During the cooking processes, the temperature
295 measured at 2 mm below the meat surface (probe 1 and 4) did not go over 120 °C. The
296 cooking conditions of every experiment are summarised in Table 2. Under these conditions,
297 the meat weight loss was affected by the duration of the marinade, as displayed in Figure 2.

298 Control samples experienced the same cooking weight loss (46-48%) regardless the meat
299 type. The minimum cooking weight loss, 18-22%, was achieved with the longest
300 marination time (>6h). When comparing cooking weight loss in the present study with an
301 earlier study using wine marinades (Busquets, Puignou, Galceran & Skog, 2006), cooking
302 weigh loss was lower with the berry marinades. This can be important when comparing the
303 effectivity of different marinades because a reduction of cooking loss, through the addition
304 of ingredients with water holding capacity, was responsible for a significant reduction on
305 the formation of PhIP and quinoxalines in burgers (Persson, Sjöholm & Skog, 2003).
306 Hence, the reduction of cooking loss, and its consequent effect on the transport of HCA
307 precursors within the meat, could play a role on decreasing the formation of HCAs in the
308 current study, besides chemical effects by the marinade components.

309 The concentrations of HCAs in unmarinated and marinated samples are reported in Table 3.
310 Among unmarinated samples (control samples), chicken, with 41 ng mutagenic HCAs/g,
311 was the most contaminated food, as compared to unmarinated camel (18 ng mutagenic
312 HCAs/g) and beef (12 ng mutagenic HCAs/g). Although there are numerous examples of

313 HCA levels in cooked chicken and beef samples reported in the literature, it is interesting to
314 know how the total levels of mutagenic HCA in unmarinated camel relate to unmarinated
315 chicken and beef cooked under the same conditions. The different concentration of HCAs
316 can be due to different levels of HCA precursors in the raw meat. For instance, Gibis &
317 Weiss (2015) confirmed that the ratio of creatin(in)e to glucose was correlated with PhIP,
318 MeIQx and harman levels in different types of cooked meat. The greater concentration of
319 PhIP in chicken, an item with low glucose concentration, was attributed to the presence of
320 certain free amino acids and creatinine (Gibis & Weiss, 2015).

321 In this study, even α -carbolines and γ -carbolines, which are traditionally reported to form at
322 300°C, were identified in chicken cooked under temperatures below 120°C (Table 3). This
323 suggests that the definition of thermal amines needs to be revised. The very sensitive
324 analysis carried out (limits of detection and recoveries in the analysis reported in
325 Supporting Information Table S1) also made possible the quantification of the Glu-P-1 and
326 Glu-P-2 in fried chicken (only). These pyridoimidazoles have been seldomly reported in the
327 literature.

328 The probable mutagens IQ and MeIQ (IARC, 1993) have been detected in the study
329 chicken samples only. These 2 quinolines were also detected, at a level with the same order
330 of magnitude, in the chicken sample (namely Shawaya) from a traditional dish prepared at a
331 Saudi restaurant (Alsohaimi, Khan, Ali & Azam, 2019). IQ and MeIQ do not form in
332 chicken exclusively as they have been detected in other matrices (e.g. fish, beef, pork and
333 goose) (Busquets, 2012; Barzegar, Kamankesh & Mohammadi, 2019). Given the high
334 toxicity of the quinolines detected in cooked meat, which have been linked to causing
335 tumours in animal studies (Sugimura, Wakabayashi, Nakagama & Nagao, 2004), the
336 consumption of fried chicken should be questioned at least for people who are at greater

337 risk of developing cancer until more is known about the link between cooked meat and
338 different types of cancer.

339 The quantification of HCAs in the 3 types of meat, with individual blueberry, raspberry and
340 strawberry marinades, which are rich in antioxidants, under conditions resembling
341 marinating in Saudi Arabian recipes, informs about the change of HCA contamination in
342 these meats caused by the berry marinades (Figure 3, Table 3). The marinades were
343 selected because that approach can be easily adopted by the public. The 3 marinades
344 affected the formation of HCAs in the 3 types of meat with a similar trend: there was a
345 strong reduction in the formation of the pyridines DMIP and PhIP; and the β -carbolines
346 harman and norharman with increased marinating time. Harman and norharman were not
347 enhanced by the marinade in this study as opposed to when common cooking recipes that
348 included multiple ingredients were used (Khan, Busquets, Naushad, Puignou, 2019).
349 Although harman levels can be correlated with glucose (Gibis & Weiss, 2015), they were
350 not increased with the application of fruit juices in this work. However, it is possible that
351 the enhancing effect of glucose on harman could be masked by the reaction caused by other
352 mechanisms.

353 Marinating for 12 and 24h was found to cause a significantly greater reduction on pyridines
354 and β -carbolines with respect to marinating for less than 6h (P 0.05). The reduction of the
355 pyridine HCAs was 91-100% and β -carbolines decreased by 40-67% with the 24h
356 marinade. Noticeably, with all 3 marinades, the concentration of quinoxalines was
357 enhanced within shorter marinating times (1h) and was reduced after 6h marination time,
358 with a 100% reduction with the 24h marination time. This trend was also observed with
359 MeIQx and 4,8-DiMeIQx when marinating with wines (Busquets, Puignou, Galceran &
360 Skog, 2006). Hence, this research shows that marinades from fruits can promote the

361 formation of quinoxalines, and that long marination time (>6h) is desirable because the
362 enhancement of quinoxalines is mitigated, probably by other chemical reactions such as the
363 capture of free radicals in the meat leading to the formation of quinoxalies. Previous works
364 demonstrated a correlation between the radical scavenging activity of the marinades and the
365 reduction of quinoxalines with time (Busquets, Puignou, Galceran & Skog, 2006; García-
366 Lomillo, Viegas, Gonzalez-SanJose & Ferreira, 2017). Future sensory analysis and
367 optimisation of the sensory properties of the prepared meat will be important to expand the
368 use of berry extracts for cooking meat.

369

370 **4. Conclusions**

371 In this study, the effect of marinating with blueberry, raspberry and strawberry on
372 commonly consumed meats has been tested under well-controlled conditions resembling
373 home cooking. Chicken was the most contaminated meat in terms of amounts of pyridines
374 and β -carbolines, with 34 ng/g and 21 ng/g respectively; followed by camel (13 and 8 ng/g)
375 and beef (7 and 6 ng/g). This study has found that marinating meat with fruit juice
376 (blueberry, raspberry and strawberry) can have a positive reduction on the formation of
377 HCAs (pyridines, carbolines and quinoxalines), especially at marinating time of at least 6h,
378 which was characterised by a 40-100% reduction in HCA. In contrast, marinades of just 1h
379 can enhance (even doubling) the formation of quinoxalines, which are potential human
380 carcinogens. The occurrence of HCAs when using 3 independent marinades was not found
381 to be dependent on the type of meat or fruit marinade. Guidelines on recommending of
382 marinating meat should emphasise on the importance of using long marination times.

383

384

385 **Acknowledgements**

386 The authors acknowledge funding from the Research and Development (R&D) Program
387 (Research Pooling Initiative), Ministry of Education, Riyadh, Saudi Arabia, (RPI-KSU).

388

389 **References:**

390 Alsohaimi, I. H., Khan, M. R., Ali, H. M., & Azam, M. (2019). Emergence of
391 mutagenic/carcinogenic heterocyclic amines in traditional Saudi chicken dishes prepared
392 from local restaurants. *Food and Chemical Toxicology*, *132*, 110677.

393 Barcelo-Barrachina, E., Moyano, E., Galceran, M. T., Lliberia, J. L., Bago, B., & Cortes,
394 M. A. (2006). Ultra-performance liquid chromatography–tandem mass spectrometry for the
395 analysis of heterocyclic amines in food. *Journal of Chromatography A*, *1125*, 195–203.

396 Barzegar, F., Kamankesh, M., & Mohammadi, A. (2019). Heterocyclic aromatic amines in
397 cooked food: A review on formation, health risk-toxicology and their analytical techniques.
398 *Food Chemistry*, *280*, 240-254.

399 Bellamri, M., & Turesky, R. J. (2019). Dietary Carcinogens and DNA adducts in prostate
400 cancer. In: Dehm S., Tindall D. (eds) Prostate cancer. Advances in experimental medicine
401 and biology, vol 1210. Springer, Cham.

402 Bellamri, M., Xiao, S., Murugan, P., Weight, C. J., & Turesky, R. J. (2018). Metabolic
403 activation of the cooked meat carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-
404 b]pyridine in human prostate. *Toxicological Sciences*, *163*, 543–556.

405 Busquets, R., Frandsen, H., Jönsson, J. A., Puignou, L., Galceran, M. T. & Skog, K. (2013).
406 Biomonitoring heterocyclic amines and metabolites in urine by liquid-phase microextraction:

407 2-amin-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP), a possible biomarker of exposure
408 to dietary PhIP. *Chemical Research in Toxicology*, 26, 233-240.

409 Busquets, R. (2012). Food borne carcinogens: a dead end? *In: Pesheva, Margarita,*
410 *Dimitrov, Martin and Stoycheva, Teodora Stefkova, (eds.) Carcinogen.* Rijeka, Croatia:
411 InTech. pp. 163-184. ISBN 9789535106586.

412 Busquets, R., Puignou, L., Galceran, M. T. & Skog, K. (2006). Effect of red wine
413 marinades on the formation of heterocyclic amines in fried chicken breast. *Journal of*
414 *Agricultural and Food Chemistry*, 54, 8376-8384.

415 Felton, J. S., Knize, M. G., Wu, R. W., Colvin, M. E., Hatch, F. T., & Malfatti, M. A.
416 (2007). Review Mutagenic potency of food-derived heterocyclic amines. *Mutation*
417 *Research*, 616, 90-94.

418 García-Lomillo, J., Viegas, O., Gonzalez-SanJose, M. L., & Ferreira, I. M. (2017). Influence
419 of red wine pomace seasoning and high-oxygen atmosphere storage on carcinogens formation
420 in barbecued beef patties. *Meat Science*, 125, 10-15.

421 Gibis, M., & Weiss, J. (2015). Impact of precursors creatine, creatinine, and glucose on the
422 formation of heterocyclic aromatic amines in grilled patties of various animal species.
423 *Journal of Food Science*, 80, C2430-C2439.

424 Gowd, V., Bao, T., & Chen, W. (2019). Antioxidant potential and phenolic profile of
425 blackberry anthocyanin extract followed by human gut microbiota fermentation. *Food*
426 *Research International*, 120, 523-533.

427 Guo, J., Villalta, P. W., Weight, C. J., Bonala, R., Johnson, F., Rosenquist, T. A., &
428 Turesky, R. J. (2018). Targeted and untargeted detection of DNA adducts of aromatic amine
429 carcinogens in human bladder by ultra-performance liquid chromatography-high-resolution
430 mass spectrometry. *Chemical Research in Toxicology*, 31, 1382-1397.

431 IARC (1993) Some naturally occurring substances: food items and constituents,
432 heterocyclic aromatic amines, and mycotoxins. IARC Monographs on the Evaluation of
433 Carcinogenic Risk of Chemicals to Humans, vol. 56, Lyon, France: International Agency for
434 Research on Cancer, pp. 165–242, 1993.

435 Khan, M. R., Busquets, R., Naushad, M. & Puignou, L. (2019). Cooking with elaborate
436 recipes can reduce the formation of mutagenic heterocyclic amines and promote comutagenic
437 amines. *Food Additives and Contaminants Part A*, 36, 385-395.

438 Khan, M. R., Busquets, R., Saurina, J., Hernández, S., & Puignou, L. (2013). Identification
439 of seafood as an important dietary source of heterocyclic amines by chemometry and
440 chromatography-mass spectrometry. *Chemical Research in Toxicology*, 26, 1014-1022.

441 Khan, M. R., Naushad, M., & Zeid, A. A. (2017). Presence of heterocyclic amine
442 carcinogens in home-cooked and fast-food camel meat burgers commonly consumed in
443 Saudi Arabia. *Scientific Reports*, 7, 1-7.

444 Manful, C. F., Vidal, N. P., Pham, T. H., Nadeem, M., Wheeler, E., Hamilton, M. C.,
445 Doody, K. M., & Thomas, R. H. (2020). Unfiltered beer based marinades reduced exposure
446 to carcinogens and suppressed conjugated fatty acid oxidation in grilled meats. *Food*
447 *Control*, 111, 107040. NTP (2004) Report on Carcinogens: 13th ed. US National Toxicology
448 Program, Department of Health and Human Services, Public Health Services, Washington
449 DC, USA, 2004. https://ntp.niehs.nih.gov/ntp/roc/content/listed_substances_508.pdf

450 Naushad, M., & Khan, M. R. *Ultra Performance Liquid Chromatography Mass*
451 *Spectrometry: Evaluation and Applications in Food Analysis*; CRC Press: Boca Raton, FL,
452 USA, 2014.

453 Oostindjer, M., Alexander, J., Amdam, G.V., Andersen, G., Bryan, N. S., Chen, D., Corpet,
454 D. E., De Smet, S., Dragsted, L. O., Haug, A., Karlsson, A. H., Kleter, G., de Kok, T. M.,

455 Kulseng, B., Milkowski, A. L., Martin, R. J., Pajari, A-M., Paulsen, J. E., Pickova, J., Rudi,
456 K., Sødning, M., Weed, D. L., & Egelanddal, B. (2014). The role of red and processed meat
457 in colorectal cancer development: a perspective. *Meat Science*, 97, 583–596.

458 Oz, F., & Kotan, G. (2016). Effects of different cooking methods and fat levels on the
459 formation of heterocyclic aromatic amines in various fishes. *Food Control*, 67, 216-224.

460 Oz, F., & Kaya, M. (2011). The inhibitory effect of black pepper on formation of
461 heterocyclic aromatic amines in high-fat meatball. *Food Control*, 22, 596-600.

462 Persson, E., Sjöholm, L., & Skog, K. (2003). Effect of high water-holding capacity on the
463 formation of heterocyclic amines in fried beefburgers. *Journal of Agricultural and Food*
464 *Chemistry*, 51, 4472-4477.

465 Rogers, L. J., Basnakian, A. G., Orloff, M. S., Ning, B., Yao-Borengasser, A., Raj, V., &
466 Kadlubar, S. (2016). 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) induces
467 gene expression changes in JAK/STAT and MAPK pathways related to inflammation,
468 diabetes and cancer. *Nutrition & Metabolism (London)*, 13, 54.

469 Skog, K.I, Johansson M.A.E., Jägerstad, M.I, Carcinogenic heterocyclic amines in model
470 systems and cooked foods: a review on formation, occurrence and intake, *Food and*
471 *Chemical Toxicology* 36 (1998) 879-896.

472 Sugimura, T., Wakabayashi, K., Nakagama, H., & Nagao, M. (2004). Heterocyclic amines:
473 Mutagens/carcinogens produced during cooking of meat and fish. *Cancer Science*, 95, 290-
474 299.

475 Vitaglione, P., & Fogliano, V. (2004). Use of antioxidants to minimize the human health
476 risk associated to mutagenic/carcinogenic heterocyclic amines in food. *Journal of*
477 *Chromatography B*, 802, 189-199.

478 Lu, F., Kuhnle, G. K., & Cheng, Q. (2017). Heterocyclic amines and polycyclic aromatic
479 hydrocarbons in commercial ready-to-eat meat products on UK market. *Food Control*, 73,
480 306-315.

481 Murkovic, M., Weber, H.J., Geiszler, S., Frohlich, K. & Pfannhauser, W. (1999).
482 Formation of the food associated carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-
483 b]pyridine (PhIP) in model systems. *Food Chemistry*, 65, 233-237.

484 Tang, Y., Kassie, F., Qian, X., Ansha, B., & Turesky, R.J. (2013). DNA adduct formation
485 of 2-amino-9H-pyrido[2,3-b]indole and 2-amino-3,4-dimethylimidazo[4,5-f]quinoline in
486 mouse liver and extrahepatic tissues during a subchronic feeding study, *Toxicological*
487 *Sciences*, 133, 248–258.

488 Turesky, R.J., & Vouros, P. (2004). Formation and analysis of heterocyclic aromatic amine
489 DNA adducts in vitro and in vivo. *Journal of Chromatography B*, 802, 155-166.

490 Zhou, L., Xie, M., Yang, F., & Liu, J. (2020). Antioxidant activity of high purity blueberry
491 anthocyanins and the effects on human intestinal microbiota. *LWT*, 117, 2020, 108621.

492
493
494
495
496
497
498
499

500 **Figure captions:**

501

502 **Figure 1:** Temperature profile obtained with type-K probes. Specifically probe 1 was
503 located the upper surface (~2 mm) of meat; probe 2 was at center of meat; probe 3, was
504 located between meat and pan surface; probe 4 was inserted within the lower layer of meat;
505 probe 5 was located at center of pan surface; and probe 6 indicated the temperature at outer
506 of the pan surface.

507

508 **Figure 2:** Meat weight loss vs. marinating time under the study conditions (n=2)

509 **Figure 3:** Variation of HCAs over marinating time in the studied meat samples.

510

511

512

513

514

515

516

517

518

519

520


521

522

523

524 **Figure 1**

525



526

527

528

529




530


531

532

533

534





539

540 **Table 1:** Cooking conditions of the meat samples processed with the marinades assayed
 541

Sample*, marinating time (h)	Sample code	Raw meat (g)	Raw meat thickness (cm)	Fruits extract (mL)	Cooking temperature (°C)	Cooking time (4 min/side)	Fried meat** (g)	Weight loss (%)
Camel ^a (control sample)	CACS	200.80	1.2	50	215-230	8.10	109.36	45.54
Camel with blueberry, (1 h)	CABL-1 h	200.74	1.1	50	215-230	8.00	125.65	37.41
Camel with raspberry, (1 h)	CARA-1 h	200.36	1.1	50	215-230	8.15	129.23	35.50
Camel with strawberry, (1 h)	SACT-1 h	200.25	1.3	50	215-230	8.15	123.65	38.25
Camel with blueberry, (6 h)	CABL-6 h	200.20	1.1	50	215-230	8.05	139.14	30.50
Camel with raspberry, (6 h)	CARA-6 h	200.35	1.1	50	215-230	8.10	142.20	29.02
Camel with strawberry, (6 h)	CAST-6 h	200.45	1.3	50	215-230	8.15	138.45	30.93
Camel with blueberry, (12 h)	CABL-12 h	200.20	1.1	50	215-230	8.00	148.89	25.63
Camel with raspberry, (12 h)	CARA-12 h	200.12	1.1	50	215-230	8.13	152.20	23.95
Camel with strawberry, (12 h)	CAST-12 h	200.32	1.3	50	215-230	8.15	147.42	26.41
Camel with blueberry, (24 h)	CABL-24 h	200.15	1.1	50	215-230	8.10	160.60	19.76
Camel with raspberry, (24 h)	CARA-24 h	200.35	1.1	50	215-230	8.10	168.23	16.03
Camel with strawberry, (24 h)	CAST-24 h	200.42	1.3	50	215-230	8.15	164.42	17.96
Beef ^b (control sample)	BECS	200.52	1.2	50	215-230	8.15	108.11	46.09
Beef with blueberry, (1 h)	BEBL-1 h	200.13	1.1	50	215-230	8.20	123.32	38.38
Beef with raspberry, (1 h)	BERA-1 h	200.42	1.3	50	215-230	8.10	125.20	37.53
Beef with strawberry, (1 h)	BEST-1 h	200.35	1.2	50	215-230	8.00	128.12	36.05
Beef with blueberry, (6 h)	BEBL-6 h	200.46	1.3	50	215-230	8.15	140.32	30.00
Beef with raspberry, (6 h)	BERA-6 h	200.42	1.2	50	215-230	8.00	135.45	32.42
Beef with strawberry, (6 h)	BEST-6 h	200.56	1.1	50	215-230	8.20	138.65	30.87
Beef with blueberry, (12 h)	BEBL-12 h	200.32	1.3	50	215-230	8.15	147.95	26.14
Beef with raspberry, (12 h)	BERA-12 h	200.85	1.3	50	215-230	8.10	152.10	24.27
Beef with strawberry, (12 h)	BEST-12 h	200.45	1.1	50	215-230	8.10	149.65	25.34
Beef with blueberry, (24 h)	BEBL-24 h	200.60	1.1	50	215-230	8.00	162.32	19.08
Beef with raspberry, (24 h)	BERA-24 h	200.78	1.2	50	215-230	8.10	161.18	19.72
Beef with strawberry, (24 h)	BEST-24 h	200.95	1.1	50	215-230	8.00	163.35	18.71
Chicken ^c (control sample)	CHCS	200.86	1.2	50	215-230	8.20	104.25	48.10
Chicken with blueberry, (1 h)	CHBL-1 h	200.65	1.1	50	215-230	8.10	117.52	41.43
Chicken with raspberry, (1 h)	CHRA-1 h	200.12	1.1	50	215-230	8.15	118.98	40.55
Chicken with strawberry, (1 h)	CHST-1 h	200.30	1.2	50	215-230	8.00	120.54	39.82
Chicken with blueberry, (6 h)	CHBL-6 h	200.25	1.3	50	215-230	8.00	141.20	29.49
Chicken with raspberry, (6 h)	CHRA-6 h	200.50	1.3	50	215-230	8.10	143.65	28.35
Chicken with strawberry, (6 h)	CHST-6 h	210.87	1.1	50	215-230	8.20	152.65	27.61
Chicken with blueberry, (12 h)	CHBL-12 h	200.45	1.2	50	215-230	8.10	149.21	25.56
Chicken with raspberry, (12 h)	CHRA-12 h	200.65	1.1	50	215-230	8.00	151.10	24.69
Chicken with strawberry, (12 h)	CHST-12 h	200.25	1.3	50	215-230	8.30	150.65	24.77
Chicken with blueberry, (24 h)	CHBL-24 h	200.50	1.2	50	215-230	8.20	162.36	19.02
Chicken with raspberry, (24 h)	CHRA-24 h	200.30	1.2	50	215-230	8.00	158.96	20.64
Chicken with strawberry, (24 h)	CHST-24 h	200.15	1.1	50	215-230	8.10	155.52	22.30

542 *Marinating temperature (4 °C); ^{a,b,c}meat cooked without fruits extract (control samples)

543

544

545

546

547

548

549

550 **Table 2.** Multiple reaction monitoring MS/MS conditions used for the quantification and
 551 confirmation of HCAs in meat samples*
 552

HCAs	Precursor ion (<i>m/z</i>) tentative assignment	Quantification		Confirmation	
		Product ion (<i>m/z</i>) tentative assignment	Collision energy (eV)	Product ion (<i>m/z</i>) tentative assignment	Collision energy (eV)
DMIP	163 [M + H] ⁺	148 [M + H - CH ₃] ⁺⁺	25	147 [M + H - CH ₃ - H] ⁺	30
PhIP	225 [M + H] ⁺	210 [M + H - CH ₃] ⁺⁺	25	183 [M + H - CH ₃ - HCN] ⁺⁺	30
Harman	183 [M + H] ⁺	115 [M + H - CH ₃ CN - HCN] ⁺	30	168 [M + H - CH ₃] ⁺⁺	30
Norharman	169 [M + H] ⁺	115 [M + H - 2HCN] ⁺	30	142 [M + H - HCN] ⁺	25
IQ	199 [M + H] ⁺	184 [M + H - CH ₃] ⁺⁺	30	157 [M + H - CH ₃ - HCN] ⁺⁺	35
MeIQ	213 [M + H] ⁺	198 [M + H - CH ₃] ⁺⁺	25	197 [M + H - CH ₃ - H] ⁺	30
MeIQx	214 [M + H] ⁺	199 [M + H - CH ₃] ⁺⁺	30	172 [M + H - CH ₃ - HCN] ⁺⁺	30
4,8-DiMeIQx	228 [M + H] ⁺	213 [M + H - CH ₃] ⁺⁺	30	187 [M + H - C ₂ NH ₃] ⁺	25
7,8-DiMeIQx	228 [M + H] ⁺	172 [M + H - CH ₃ - C ₂ NH ₃] ⁺⁺	35	213 [M + H - NH ₃] ⁺⁺	25
4,7,8-TriMeIQx (IS)	242 [M + H] ⁺	227 [M + H - CH ₃] ⁺⁺	25	201 [M + H - C ₂ NH ₃] ⁺	30
Glu-P-1	199 [M + H] ⁺	172 [M + H - HCN] ⁺	25	184 [M + H - CH ₃] ⁺⁺	25
Glu-P-2	185 [M + H] ⁺	158 [M + H - HCN] ⁺	25	131 [M + H - HCN - HCN] ⁺	30
AαC	184 [M + H] ⁺	167 [M + H - NH ₃] ⁺	25	140 [M + H - NH ₃ - HCN] ⁺	30
MeAαC	198 [M + H] ⁺	181 [M + H - NH ₃] ⁺	25	154 [M + H - NH ₃ - HCN] ⁺	30
Trp-P-1	212 [M + H] ⁺	195 [M + H - NH ₃] ⁺	25	168 [M + H - NH ₃ - HCN] ⁺	30
Trp-P-2	198 [M + H] ⁺	154 [M + H - NH ₃ - HCN] ⁺	30	181 [M + H - NH ₃] ⁺	25

553 *System dwell time was 0.025 s in all studied compounds; IS, internal standard
 554
 555

556

557

558

559

560

561

Table 3. HCAs identified in thermally processed camel, beef and chicken meat samples marinated with highly antioxidant fruits. The acronyms CA(camel), BE (beef), CH (chicken), BL(blueberry);RA(raspberry);ST (strawberry) are used.

Sample code	DMIP (ng/g) ± sd	PhIP (ng/g) ± sd	Harman (ng/g) ± sd	Norharman (ng/g) ± sd	IQ (ng/g) ± sd	MeIQ (ng/g) ± sd	MeIQx (ng/g) ± sd	4,8-DiMeIQx (ng/g) ± sd	7,8- DiMeIQx (ng/g) ± sd	Glu-P-1 (ng/g) ± sd	Glu-P-2 (ng/g) ± sd	AαC (ng/g) ± sd	MeAαC (ng/g) ± sd	Trp-P-1 (ng/g) ± sd	Trp-P-2 (ng/g) ± sd
CACS ^a	4.23 ± 0.31	8.65 ± 0.53	2.36 ± 0.16	5.45 ± 0.34	nd	nd	2.85 ± 0.15	1.82 ± 0.10	0.84 ± 0.03	nd	nd	nd	nd	0.03 ± 0.002	0.03 ± 0.002
CABL-1 h	1.82 ± 0.08	4.36 ± 0.04	1.82 ± 0.07	4.68 ± 0.26	nd	nd	3.62 ± 0.12	2.54 ± 0.13	1.06 ± 0.02	nd	nd	nd	nd	0.01 ± 0.001	0.01 ± 0.001
CARA-1 h	2.68 ± 0.12	5.45 ± 0.06	1.94 ± 0.09	4.82 ± 0.28	nd	nd	3.38 ± 0.14	2.63 ± 0.14	1.21 ± 0.04	nd	nd	nd	nd	0.01 ± 0.001	0.01 ± 0.001
CAST-1 h	2.42 ± 0.11	4.67 ± 0.04	1.87 ± 0.08	4.72 ± 0.26	nd	nd	3.74 ± 0.12	2.86 ± 0.13	1.06 ± 0.05	nd	nd	nd	nd	nd	nd
CABL-6 h	1.62 ± 0.05	4.03 ± 0.04	1.62 ± 0.05	4.12 ± 0.21	nd	nd	1.52 ± 0.12	1.26 ± 0.10	0.01 ± 0.001	nd	nd	nd	nd	nd	nd
CARA-6 h	2.41 ± 0.12	4.85 ± 0.05	1.74 ± 0.06	4.19 ± 0.23	nd	nd	1.75 ± 0.13	1.45 ± 0.11	0.01 ± 0.001	nd	nd	nd	nd	nd	nd
CAST-6 h	2.12 ± 0.10	4.21 ± 0.03	1.68 ± 0.06	4.15 ± 0.24	nd	nd	1.63 ± 0.12	1.34 ± 0.11	0.01 ± 0.001	nd	nd	nd	nd	nd	nd
CABL-12 h	1.26 ± 0.03	2.35 ± 0.02	1.48 ± 0.04	3.85 ± 0.18	nd	nd	0.87 ± 0.05	0.56 ± 0.03	nd	nd	nd	nd	nd	nd	nd
CARA-12 h	1.42 ± 0.05	2.67 ± 0.02	1.53 ± 0.05	3.89 ± 0.20	nd	nd	0.99 ± 0.08	0.69 ± 0.04	nd	nd	nd	nd	nd	nd	nd
CAST-12 h	1.38 ± 0.04	2.53 ± 0.02	1.61 ± 0.04	3.87 ± 0.20	nd	nd	0.92 ± 0.08	0.63 ± 0.04	nd	nd	nd	nd	nd	nd	nd
CABL-24 h	nd	0.66 ± 0.04	1.36 ± 0.03	3.21 ± 0.16	nd	nd	nd	nq	nd	nd	nd	nd	nd	nd	nd
CARA-24 h	nd	0.74 ± 0.04	1.38 ± 0.03	3.26 ± 0.13	nd	nd	nd	nq	nd	nd	nd	nd	nd	nd	nd
CAST-24 h	nd	0.68 ± 0.04	1.42 ± 0.02	3.24 ± 0.17	nd	nd	nd	nq	nd	nd	nd	nd	nd	nd	nd
BECSf ^b	2.34 ± 0.13	4.72 ± 0.34	2.25 ± 0.11	3.61 ± 0.18	nd	nd	2.13 ± 0.13	1.74 ± 0.12	0.67 ± 0.05	nd	nd	nd	nd	nd	nd
BEBL-1 h	1.95 ± 0.10	3.35 ± 0.23	1.95 ± 0.10	2.61 ± 0.11	nd	nd	3.46 ± 0.17	2.12 ± 0.11	0.78 ± 0.04	nd	nd	nd	nd	nd	nd
BERA-1 h	2.23 ± 0.12	3.65 ± 0.43	1.98 ± 0.10	2.68 ± 0.12	nd	nd	3.40 ± 0.16	2.33 ± 0.12	0.82 ± 0.03	nd	nd	nd	nd	nd	nd
BEST-1 h	2.05 ± 0.12	3.21 ± 0.53	2.03 ± 0.13	2.85 ± 0.12	nd	nd	3.43 ± 0.16	2.02 ± 0.11	0.46 ± 0.06	nd	nd	nd	nd	nd	nd
BEBL-6 h	1.82 ± 0.10	3.25 ± 0.63	1.86 ± 0.12	2.54 ± 0.10	nd	nd	0.17 ± 0.01	0.01 ± 0.001	nq	nd	nd	nd	nd	nd	nd
BERA-6 h	2.01 ± 0.12	3.32 ± 0.43	1.85 ± 0.12	2.61 ± 0.13	nd	nd	0.19 ± 0.01	0.01 ± 0.001	nq	nd	nd	nd	nd	nd	nd
BEST-6 h	1.98 ± 0.12	2.86 ± 0.02	1.76 ± 0.10	2.65 ± 0.13	nd	nd	0.28 ± 0.02	0.01 ± 0.001	nq	nd	nd	nd	nd	nd	nd
BEBL-12 h	0.54 ± 0.02	0.72 ± 0.04	1.32 ± 0.08	1.48 ± 0.06	nd	nd	nq	nd	nd	nd	nd	nd	nd	nd	nd
BERA-12 h	0.46 ± 0.01	0.64 ± 0.03	1.45 ± 0.08	1.62 ± 0.07	nd	nd	nq	nd	nd	nd	nd	nd	nd	nd	nd
BEST-12 h	0.53 ± 0.01	0.68 ± 0.04	1.38 ± 0.07	1.56 ± 0.06	nd	nd	nq	nd	nd	nd	nd	nd	nd	nd	nd
BEBL-24 h	nd	0.01 ± 0.001	1.23 ± 0.05	1.42 ± 0.04	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
BERA-24 h	nd	0.02 ± 0.002	1.35 ± 0.05	1.56 ± 0.03	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
BEST-24 h	nd	0.01 ± 0.001	1.28 ± 0.04	1.53 ± 0.03	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
CHCS ^a	8.82 ± 0.51	24.95 ± 2.44	5.87 ± 0.55	14.62 ± 1.82	0.05 ± 0.003	0.23 ± 0.03	3.55 ± 0.18	2.84 ± 0.14	0.14 ± 0.006	0.04 ± 0.003	0.02 ± 0.001	0.06 ± 0.004	0.14 ± 0.02	0.03 ± 0.002	0.03 ± 0.002
CHBL-1 h	6.23 ± 0.35	20.55 ± 2.20	4.14 ± 0.46	12.61 ± 1.64	0.03 ± 0.002	0.17 ± 0.02	4.87 ± 0.16	3.95 ± 0.12	0.29 ± 0.06	0.03 ± 0.002	nq	0.04 ± 0.003	0.08 ± 0.005	0.01 ± 0.001	0.01 ± 0.001
CHRA-1 h	7.21 ± 0.45	21.12 ± 2.20	5.22 ± 0.54	12.68 ± 1.58	0.04 ± 0.003	0.19 ± 0.02	4.03 ± 0.13	3.77 ± 0.15	0.22 ± 0.08	0.02 ± 0.001	nq	0.03 ± 0.002	0.03 ± 0.002	0.02 ± 0.001	0.02 ± 0.002
CHST-1 h	6.72 ± 0.40	20.87 ± 2.10	4.35 ± 0.32	12.85 ± 1.87	0.03 ± 0.002	0.18 ± 0.02	4.94 ± 0.12	3.82 ± 0.16	0.26 ± 0.05	0.02 ± 0.001	nq	0.03 ± 0.002	0.02 ± 0.001	0.01 ± 0.001	0.01 ± 0.001
CHBL-6 h	5.21 ± 0.36	18.65 ± 2.00	3.86 ± 0.25	12.54 ± 1.71	0.02 ± 0.001	0.13 ± 0.01	2.17 ± 0.12	1.78 ± 0.14	0.66 ± 0.05	nq	nd	nq	0.06 ± 0.004	nd	nd
CHRA-6 h	5.65 ± 0.41	19.44 ± 2.20	3.85 ± 0.26	12.52 ± 1.67	0.03 ± 0.002	0.16 ± 0.02	2.19 ± 0.12	1.84 ± 0.14	0.71 ± 0.06	nq	nd	0.02 ± 0.001	0.04 ± 0.003	nd	nd
CHST-6 h	5.33 ± 0.40	18.13 ± 2.22	3.76 ± 0.16	12.54 ± 1.89	0.02 ± 0.001	0.14 ± 0.01	2.28 ± 0.14	1.82 ± 0.13	0.69 ± 0.07	nq	nd	0.02 ± 0.001	0.03 ± 0.003	nd	nd
CHBL-12 h	1.65 ± 0.10	6.69 ± 0.32	2.46 ± 0.14	10.42 ± 1.30	nd	nd	1.18 ± 0.05	0.56 ± 0.04	nq	nd	nd	nd	0.01 ± 0.001	nd	nd
CHRA-12 h	2.24 ± 0.02	7.84 ± 0.34	2.58 ± 0.32	10.35 ± 1.25	nd	nd	1.42 ± 0.06	0.84 ± 0.06	nq	nd	nd	nd	nd	nd	nd
CHST-12 h	1.74 ± 0.11	6.36 ± 0.34	2.49 ± 0.32	10.46 ± 1.24	nd	nd	1.26 ± 0.04	0.71 ± 0.05	nq	nd	nd	nd	nq	nd	nd
CHBL-24 h	0.56 ± 0.03	1.33 ± 0.06	1.95 ± 0.13	7.89 ± 0.51	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
CHRA-24 h	0.71 ± 0.04	2.15 ± 0.12	2.33 ± 0.10	8.72 ± 0.56	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
CHST-24 h	0.68 ± 0.04	1.41 ± 0.05	2.11 ± 0.12	7.64 ± 0.49	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd

^{a,b,c}Cooked without addition of fruit juice (control samples); sd, standard deviation (n = 3), obtained from addition standard calibration curve; nq: below quantification limit; nd, not detected

Supporting information

Blueberry, Raspberry, and Strawberry Extracts Reduce the Formation of Carcinogenic Heterocyclic Amines in Fried Camel, Beef and Chicken meats.

Mohammad Rizwan Khan¹, Rosa Busquets*², Mohammad Azam¹

¹Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

²School of Life Sciences, Pharmacy and Chemistry, Kingston University London, Penrhyn road, KT1 2EE, Kingston upon Thames, United Kingdom

Figure S1: Structures and abbreviations of the study HCAs.




Table S1. HCAs limit of detection (LOD) and recovery (R) in cooked camel, beef and chicken meat

HCAs	CACs ^a		CABL-1 h		CARA-1 h		SACT-1 h		BECS ^b		BEBL-1 h		BERA-1 h		BEST-1 h		CHCS ^c		CHBL-1 h		CHRA-1 h		CHST-1 h	
	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %	LOD, ng/g	R, %
DMIP	0.03	84	0.04	77	0.03	79	0.04	81	0.03	83	0.05	78	0.04	77	0.04	80	0.01	86	0.03	81	0.02	78	0.02	82
PhIP	0.01	45	0.03	38	0.02	40	0.02	42	0.01	48	0.02	43	0.03	42	0.02	44	0.01	52	0.02	44	0.02	43	0.03	41
Harman	0.01	88	0.02	84	0.03	82	0.02	86	0.01	89	0.02	83	0.02	85	0.03	83	0.01	90	0.02	87	0.02	86	0.02	87
Norharman	0.01	86	0.02	83	0.02	84	0.02	85	0.01	87	0.03	80	0.02	81	0.03	80	0.01	89	0.02	85	0.02	86	0.02	85
IQ	0.02	76	0.04	72	0.03	70	0.03	73	0.02	77	0.03	73	0.04	72	0.02	74	0.01	78	0.02	75	0.03	74	0.03	76
MeIQ	0.02	27	0.03	24	0.02	23	0.03	24	0.02	26	0.02	23	0.03	22	0.01	29	0.02	24	0.03	25	0.03	24	0.03	25
MeIQx	0.02	38	0.03	35	0.04	32	0.03	34	0.02	36	0.04	31	0.03	32	0.04	31	0.01	41	0.02	36	0.02	35	0.02	37
4,8-DiMeIQx	0.01	54	0.03	48	0.02	46	0.03	48	0.02	50	0.03	48	0.02	46	0.03	45	0.01	56	0.02	53	0.03	51	0.03	48
7,8- DiMeIQx	0.02	50	0.03	48	0.03	45	0.02	43	0.02	48	0.03	43	0.03	45	0.03	46	0.01	52	0.02	53	0.02	50	0.02	49
Glu-P-1	0.03	36	0.04	31	0.04	29	0.03	27	0.03	33	0.04	28	0.03	30	0.02	37	0.02	39	0.03	35	0.03	36	0.03	35
Glu-P-2	0.03	40	0.03	37	0.03	35	0.04	35	0.03	38	0.04	36	0.04	35	0.02	43	0.02	42	0.04	36	0.03	38	0.03	37
A α C	0.02	26	0.03	23	0.03	22	0.04	21	0.02	24	0.03	22	0.03	20	0.03	19	0.01	28	0.03	24	0.02	25	0.02	26
MeA α C	0.01	78	0.02	75	0.02	74	0.03	72	0.01	76	0.02	72	0.03	71	0.02	73	0.01	80	0.02	77	0.03	76	0.02	75
Trp-P-1	0.02	45	0.03	39	0.04	38	0.03	35	0.02	43	0.03	37	0.03	36	0.02	38	0.01	48	0.02	42	0.02	42	0.02	43
Trp-P-2	0.01	41	0.02	38	0.03	35	0.03	36	0.02	38	0.03	33	0.03	32	0.03	31	0.01	43	0.02	38	0.03	37	0.03	38

^{a,b,c}Fried without the addition of fruit extract (control samples); LOD was estimated as the concentration with a signal-to-noise ratio of 3:1; CACS, camel (control sample); CABL, camel with blueberry; CARA, camel with raspberry; SACT, camel with strawberry; BECS, beef (control sample); BEBL, beef with blueberry; BERA, beef with raspberry; BEST, beef with strawberry; CHCS, chicken (control sample); CHBL, chicken with blueberry; CHRA, chicken with raspberry; CHST, chicken with strawberry

Credit Author Statement Mohammad Rizwan Khan: Conceptualization, Methodology, Investigation, Funding acquisition. Rosa

Busquets: Data curation, Validation, Writing- Original draft preparation, Supervision. Mohammad Azam: Formal analysis,

Investigation.