This is an Accepted Manuscript of an article published in *BMJ Quality & Safety* on 25th February 2025, available at: https://doi.org/10.1136/bmjqs-2024-017560

- 1 A scoping review identifying interventions that have been tested to
- 2 optimise the experience of people from ethnic minority groups
 - receiving systemic anti-cancer therapy (SACT)
- 4

5 Abstract

6 Background

Disparities have been identified in many aspects of the cancer care pathway for people from minority
ethnic groups (MEG). Adherence to systemic anti-cancer therapies (SACT) has been shown to impact
morbidity and mortality and therefore unequitable experiences can have a detrimental effect on
outcomes.

11 Objectives

To identify interventions that focused on improving the experiences and clinical outcomes in peoplefrom MEG receiving SACT treatments.

14 Methods

- 15 A scoping review was conducted according to the Arksey and O'Malley's methodological framework
- 16 to map the available literature. A comprehensive search was performed using three electronic
- 17 databases (Medline, Embase and Cinahl). Standard scoping review methodology following PRISMA
- 18 guidelines was used. Studies were included that assessed interventions to improve MEG patients
- 19 experience with SACT. Study types included in the review were evaluation studies, randomised/non-
- 20 randomised controlled trials and all observational studies. Exclusion criteria were applied to studies
- 21 including opinion pieces, literature and systematic reviews, non-English studies, conference abstracts
- and studies that were not describing an intervention. Independent duplicate screening, study
- 23 selection, data extraction and quality assessment was undertaken. Results of the studies were
- 24 synthesised using a published equity framework.

25 Results

- 26 Searches yielded 1,356 articles. Nine studies were included after exclusion criteria were applied.
- 27 Studies described six digital, two in person and one hybrid interventions employing different
- 28 research methodologies, ranging from randomised controlled trials (RCT's), feasibility study and
- 29 mixed methods studies. The majority of interventions in this study were delivered remotely, using
- 30 digital platforms such as websites, recorded educational training materials as well as social media.
- 31 These interventions were conducted in the USA and primarily targeted early breast cancer patients
- 32 from African American backgrounds.

33 Conclusions

- 34 This scoping review showed that there has been a very small number of studies investigating
- 35 interventions to optimise SACT treatment experiences in people from MEG. We found evidence of
- 36 interventions incorporating the equity domains that reported improved patient engagement and
- 37 experience. This new knowledge will help to implement future SACT interventions, addressing health
- 38 inequities across the cancer continuum.
- 39

Key messages

2 What is already known on this topic: Despite decades of global efforts to reduce cancer disparities,

3 patients with cancer continue to face inequalities throughout their cancer journey, including access

- 4 to and treatment with Systemic Anti-Cancer Therapy (SACT).
- 5 What this study adds: Our review identified interventions that were explicitly designed to
- 6 incorporate equity domains. Interventions incorporating these domains reported improvements in
- 7 patient experience however the number of studies was small and they used different evaluation
- 8 designs.

9 How this study might affect research, practice or policy: All the interventions in this study were

10 conducted in the USA, highlighting an urgent need for future research based in the UK to ensure that

11 patients receiving Systemic Anti-Cancer Therapy (SACT) have access to equitable cancer care. Future

12 studies should prioritise exploring psychosocial and cultural influences to inform the development of

13 effective and equitable interventions tailored to the needs of diverse patient populations.

14

15 Introduction

16 Cancer is the leading cause of mortality worldwide and over the past two decades, innovations in drug

17 treatment have improved overall survival [1]. Improvements are not equally applicable to all patients,

18 with significant differences in cancer mortality rates between people from minority ethnic groups

[MEG] [2]. Although considerable progress has been made in the last decade describing cancer health
 disparities amongst people from MEG, inequalities still exist, which was further highlighted during

21 COVID-19 pandemic. [3]

22 The specific health barriers that people from MEG are still facing today are multifactorial and are 23 influenced by patients' demographics (age, gender, insurance, social class, race, and geography), 24 language and acculturation, attitudes, and family and cultural contexts. [4] Persistent lack of resources 25 to protect and improve health, influences how people from MEG perceive healthcare services and are 26 underpinned by intermediary, social and structural determinants of health. [5] At present in the United 27 Kingdom (UK) there is limited research evidence to indicate what types of intervention would prove 28 most effective among MEG and marked differences among the UK population exist mainly due to lack 29 of preventative care in general. [6]

With the World Health Organisation's declaration, post COVID-19 public health emergency, attention has shifted to the prevention and treatment of other communicable diseases as well as noncommunicable diseases, and the critical contributions of medicines globally. Cancer incidence is expected to rise significantly through 2050, particularly in lower-income countries, with an increase of over 12 million new cases annually. The growing disease burden underscores the urgent need for new

35 treatments to enhance survival and quality of life. [7]

36 Drug treatments for cancer are available in many forms; oral, intravenous, sub-cutaneous and intra-37 thecal [1]. There has been a rise in the use of oral therapies in the last 10 years requiring administration 38 and correct dosing to be managed by the person with cancer, over long periods of time. Oral therapies 39 are convenient for the patient as they reduce frequent visits to cancer centres to receive intravenous 40 drugs and enables patients to self-manage their treatment. However, patient adherence to oral 41 anticancer drugs is an emerging issue in modern oncology. A systematic review by Greer et.al. reported 42 that patients' adherence to oral SACT can be as low as 46%. [8] A further study investigated rates of 43 non-adherence between people from a White British background and ethnic minority breast cancer

- 1 survivors finding that women from minority ethnic groups had a significantly higher risk of non-
- 2 adherence than women who were from a White British background (odds ratio = 1.50, p = 0.03). [9]
- 3 Patients are taking medications for prolonged periods of time for certain cancers and are expected to
- 4 notice and report serious side effects to their medical team. Without tailored patient education, it is
- 5 difficult to achieve maximum benefit for the individual patient and consequently for the whole health
- 6 system.
- 7 Statistics in the UK show that more than a third (36%) of all cancer cases are diagnosed in people
- 8 aged 75 and over, and with age the number of comorbidities requiring medication also rises [10,11].
- 9 Patients on multiple medications may struggle to adhere to Systemic Anti-Cancer Therapy (SACT)
- 10 treatment regimen and there are many factors influencing non-adherence to oral anti-cancer drugs,
- 11 such as socioeconomic-related factors, healthcare system-related factors, patient-related factors,
- 12 disease-related factors, and therapy-related factors [12]. Non-adherence to medications negatively
- 13 affects efficacy, safety and costs of therapies. Evidence suggests that there is a social gradient to non-
- 14 adherence, as belonging to an ethnic minority might have a negative impact on adherence to
- 15 medications, which is closely related to the level of health literary. [13]
- 16 Health literacy is defined as "the degree to which individuals can obtain, process and understand the 17 basic health information to make appropriate health decisions". [14] It is known that high health 18 literacy is associated with increased patient engagement, better self-management and essentially 19 improved health outcomes [12]. There is evidence to suggest that healthcare professionals should be 20 addressing unconscious biases when identifying patients with low literacy levels rather than targeting 21 knowledge deficit .[15] Implicit biases are unconscious attitudes and beliefs, that influence people's 22 behaviour and interracial interactions often may produce mistrust in healthcare.[16] While there are 23 increasing efforts to explore how treatments can be improved for people from MEG in Western 24 societies, a holistic global review of this pressing matter does not yet exist. This may, in part, be due to 25 the tendency among researchers to focus on specific countries, diseases and impacted communities 26 [17]. The intersectionality of ethnicity with other factors such as advanced age, health and digital 27 literacy compound the inequity.
- The present review draws together available evidence on the types of interventions used to optimise cancer patients from MEG treatment with systemic anticancer therapy (SACT). The findings from this
- 30 review will inform the design of potential interventions, to improve adherence and subsequent
- 31 outcomes to achieve equitable cancer care.
- The review was guided by three research questions that supported the aim of identifying interventions that have been used to optimise SACT treatment in patients of different ethnicities.
- 34 1. How were interventions conducted in their chosen patient population?
- 35 2. Have these interventions improved patient experience, safety and efficacy of treatments?
- 36 3. What were the key equity dimensions related to these interventions?
- 37
- 38 Methods
- 39 Protocol and registration
- 40 This scoping review followed the Arskey and O'Malley methodological framework for scoping
- 41 reviews [18] and the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension
- 42 for Scoping Reviews (PRISMA-ScR) checklist criteria [19] [See Appendix 1]. The study protocol was
- 43 registered on Open Science Framework (registration DOI 10.17605/OSF.IO/KFS7H).
- 44

1 <u>Search strategy</u>

- 2 The population, intervention, comparison, outcome (PICO) framework [20] was used to develop the
- 3 search strategy. Following this, a comprehensive search was performed from database inception until
- 4 June 7th, 2023 with the following databases: EMBASE (Ovid), MEDLINE (Ovid) and Cumulative Index
- 5 to Nursing and Allied Health Literature (CINAHL). The search strategies used MESH terms and key
- 6 words outlined in Appendix 2. The reference lists of the 7 selected publications were searched for
- 7 additional sources and additional 2 publications were identified. A combined initial screening of title
- 8 and abstract for eligibility was then conducted using a priori study protocol. Following the initial
- 9 screening at the title and abstract level, a second researcher (author LS) verified 10% of exclusions
- 10 against the inclusion criteria. Subsequently, the remaining publications that met the inclusion criteria
- 11 underwent full-text screening for eligibility. To ensure improved accuracy, 100% of included and 5%
- 12 of excluded documents were checked by author LS, and discrepancies were resolved via consensus.
- 13 The search was re-run on the 6th of August 2024 with the following databases: EMBASE (Ovid) and
- 14 MEDLINE (Ovid); no new studies were identified.
- 15

16 Eligibility criteria

- 17 Eligibility criteria are summarised in Table 1. The population of this study was defined as people from
- 18 MEG, classified based on a combination of social, cultural or linguistic characteristics that would
- 19 distinguish them from the majority of the population in the United Kingdom (UK). All cancer types
- 20 were included in the study and the types of interventions that were considered ranged from
- 21 educational programs, behavioural interventions, changes in healthcare delivery models to
- 22 pharmacological adjustments. The delivery of interventions could be provided by any type of
- 23 healthcare provider and the setting for interventions was not limited to clinical, online or
- 24 community-based services. The outcome of patient experiences was defined by numbers of health
- 25 interactions or surveys capturing satisfaction within interventions, that influenced patient
- 26 perceptions to various outcomes as defined in Table 1.

Eligibility criteria	Population	Intervention	Comparison	Outcome
criteria Inclusion criteria	Adult (over 18 y) People from MEG Receiving SACT	Studies including interventions to better manage systemic anti- cancer therapies (SACT) Studies exploring implementation of interventions Study types included in the review were evaluation studies, randomised/non-randomised controlled trials and all observational studies.	Usual care/No intervention	Adherence and compliance to therapy (SACT) defined by either patient following prescribed treatment regimen or treatment completion. Quality of life (measured using QOL assessment tools) Adverse events to treatments Symptom management Self-efficacy Barriers and limitations to implementation of interventions Patient experiences of intervention delivery

27 [Table 1] PICO search strategy

				Healthcare utilization, survival rates, cost- effectiveness, disparities in care
Exclusion criteria	Patients under 18 years old	Studies including opinion pieces, literature and systematic reviews, non-English studies, conference abstracts Studies not describing an intervention Not retrievable studies	N/A	N/A

- 2 Key characteristics of the interventions described the following parameters in these studies: author,
- 3 year, country, intervention type, objectives, methods, sample demographics, outcomes, dimensions
- 4 of accessibility to interventions, quality assessment of interventions.

5 <u>Selection of sources of evidence</u>

- 6 The RefWorks bibliographic software package was used to manage all the references. Potential biases
- 7 were considered during the selection process. The inclusion of a specific population in the study can
- 8 have a dramatic impact on the conclusions for the effectiveness of a treatment. [21] It is known that
- 9 racial disparities among cancer patients are a widespread phenomenon affecting health outcomes of
- 10 this group of patients, so by selecting this population could introduce a spectrum bias and
- 11 publication bias. To avoid these biases two researchers conducted the selection process, including
- 12 studies with positive and negative research findings. The reviewers verified that the studies met the
- 13 eligibility criteria and addressed our research question.

14 Data charting process

- Studies that met inclusion criteria were summarised qualitatively, and the summary of the resultsreported according to the Arskey and O'Malley guidelines [18].
- 17 Two Excel data extraction forms were used to organise the review process:
- 18 1. Data were extracted from articles and charted using the unique chart adapted from the 19 Template of intervention description and Republication checklist [Appendix 3] [22].
- Template of intervention description and Republication checklist [Appendix 3] [22].
 The methodological quality of the empirical studies was critically appraised by two
 - The methodological quality of the empirical studies was critically appraised by two researchers using the Mixed Methods Appraisal Tool (MMAT) [Appendix 4] [23].
- 22 Participant and design characteristics of the studies are summarised in table 2. Following extraction

23 of the data, studies were summarised in table 3. and included the following details: author, year,

24 country, intervention, methods, sample size, study population, cancer type and specific study

25 outcomes.

21

26 Data synthesis and reporting

- 27 Data were synthesized using the equity framework adapted from Sayani et al. [5]. To assess
- 28 accessibility dimensions of the interventions there were five parameters used in the process:
- 29 approachability, acceptability, availability, affordability, and appropriateness of the studies.
- 30 Interventions were mapped against the result section of the framework using five dimensions and
- 31 the summary is provided in figure 1. The features listed under each dimension in Figure 1 were based
- 32 on Sayani et.al work, as well as clinical practice and early conversations with the patients and
- 33 partners in this study.

- 1 Only the data relevant to SACT was extracted from the original equity framework, which was
- 2 designed to assess effectiveness of interventions in accessing lung cancer screening amongst
- 3 targeted priority populations. Five parameters that were used to analyse the intervention
- 4 descriptions and the definition of each is provided as follows:
- 1) Approachability parameter included information exploring the ability of participants to perceive
 risk about the available support services, recruitment process and personalised risk assessment.
- 7 2) Acceptability parameter assessed the ability to seek information through social and cultural
- 8 factors determining participation in interventions, such as the use of translator services, sociocultural
- 9 training for staff as well as translating materials into native languages.
- 3) Availability assessed the ability to reach wider communities for participation in the intervention,
 including ability to use technology and transportation considerations.
- 4) Affordability assessed any costs associated with SACT treatment and whether indirect costs wereconsidered when designing an intervention.
- 14 5)Appropriateness parameter was used to assess patient's engagement and empowerment to make
- 15 informed decisions during interventions.
- 16

17 Results

- 18 <u>Search results</u>
- 19 The initial search of the databases yielded 1356 articles of which 106 were taken to abstract review
- 20 following removal of duplicates (n=41) (Figure 2). 1209 articles were removed after title screen,
- 21 which was performed by researcher (JM) with a 10% validation performed by researcher (LS). Full
- 22 abstract review was performed by two researchers (JM, LS). 41 articles were excluded after the
- abstract screening stage leaving 65 articles for full review. Following the application of inclusion and
- exclusion criteria by both authors, seven studies were included in analysis and quality assessment.
- 25 An additional, two studies were identified through citation searches, resulting in a total of nine
- 26 studies included in this scoping review. Mixed Methods Appraisal Tool (MMAT) was used to assess
- the overall quality of the studies and the overall score provided in table 3. The detailed analysis of
- the quality of studies are summarised in Appendix 4 [23]. To explore types of interventions the
- 29 template of intervention description and Republication checklist was adapted and used to analyse
- 30 data [Appendix 3] [22].
- 31 32
 - Table 2. Participant and design characteristics of studies in the scoping review (n = 9).
- 33

Study characteristics	Number (n)
Cancer type:	
Breast	n=7
Multiple	n=1
Breast and Lung	n=1
Sample size (range)	n=24-1442
Participant age (range)	n=45-63
Study design:	
RCT	n=3
NRCT	n=1
Mixed Methods study	n=1
Randomised pilot study	n=3

Non-Randomised pilot study	n=1
Intervention type:	
In person	n=2
Digital	n=6
Hybrid (Digital/In person)	n=1
Stage of cancer:	
1	n=5
II	n=0
111	n=0
IV	n=0
Any stage	n=4
Ethnicity of participants:	
African Americans	n=7
Asian Americans	n=1
Hispanic/Latino	n=1

2 Description of interventions designed to optimise experience with SACT for people from MEG

3 Key characteristics of the studies were summarised in table 2. There was much heterogeneity within

4 the study populations and outcomes for the nine included studies. Interestingly, all the studies were

5 conducted in the United States of America (USA) with a wide range of sample sizes, including studies

6 with 24 participants and larger studies recruiting 1442. Studies employed different designs to deliver

7 interventions and were carried out by various healthcare professionals, often race matched (n=5)

8 with the participants in the studies. The length of the studies ranged from a single two-hour

9 workshop intervention (n=1) to studies supporting patients through care navigation and highlighting

10 missed appointments to providers over a 5-year period (n=1). Seven different categories of outcome

11 variables were examined across the 9 studies: medication adherence (n=2), health literacy (n=1), 12 superior of two two ends of the efficiency (n=2) superior of life (n=2) superior of life (n=2).

12 completion of treatment (n=1), self-efficacy (n=2), quality of life (n=2), survivorship experience (n=1),

13 feasibility of an intervention (n=2).

14 Digital interventions versus in-person and hybrid interventions

- 15 Of the nine studies, six interventions were conducted digitally, two were delivered in person and
- 16 another study incorporated mixed mode of delivery. (Appendix 3) The majority of interventions used
- 17 technology to optimise patients' experiences with SACT. There were only two studies that assessed
- 18 quality of life of participants and both of them showed no effect post intervention [24,25]. The rest
- 19 of the four digital studies had positive outcomes, including improved completion rates of treatment
- 20 [26] and high satisfaction with the programme's content on the websites [27,28]. Randomised pilot
- 21 study conducted by Perez et. al reported moderate-to-high levels of positive emotional reactions to
- 22 stories and identification with storytellers.[29] Digital interventions used different approaches to
- reach participants in their studies. Two of the studies designed culturally tailored virtual
- 24 programmes, available on their websites [27,28], three studies employed interactive videos showing
- race-matched survivor stories [24,25, 29] and further digital study used electronic record system that
- 26 would flag up missed appointments or unmet milestones in patient's treatment. [30]
- 27 Out of two in-person delivered interventions, one did not show an effect on health literacy,
- 28 medication adherence or self-efficacy from pre-test to post intervention. [30]. Another in person
- 29 delivered study was a feasibility trial and reported increased self-efficacy in communicating with
- 30 providers and self-efficacy in making treatment decisions. [31]
- 31 An intervention conducted by Rosenzweig et.al. incorporated supportive video messages from the
- 32 African American community as well as in person training in their design. This hybrid study reported

1 a positive effect of a psychoeducational session with race-matched interventionist delivering training

- 2 sessions. [32]
- 3
- -
- 4
- 5

6 Equity assessment of interventions designed to optimise experience with SACT for people from MEG

- 7 People from MEG were the focus population of the nine studies, where participants were mainly
- 8 recruited from urban locations (n=7), 17 community oncology practices (n=1) and online (n=1). All of
- 9 the studies incorporated culturally targeted materials in their design, ranging from guidebooks, print
- 10 materials and decision-making models. There were several similarities across the studies, such as
- 11 identification of the need to race match the recruiter with study participants [30,32], recognising the
- 12 importance of translating educational materials into native languages [24,28] as well as providing
- 13 staff with racial equity training [25,26,29].
- 14 Interventions were analysed using the equity framework as reported by Sayani et al. [5],
- 15 incorporating different accessibility dimensions such as approachability, acceptability, availability,
- 16 affordability, and appropriateness in people from MEG. (Table 4) All the interventions suggested that
- 17 approachability to SACT could be enhanced through raising awareness and recruiting participants
- 18 from different geographic locations, including both online and offline strategies.
- 19 Acceptability of interventions was reported in all but one study [30], which did not incorporate
- 20 neither cultural staff training, translation of materials into other languages or tailoring intervention
- to personal and cultural values of participants. Only four interventions considered the availability of
- 22 SACT across people from MEG, as they did not rely on participants ability to use technology, and
- 23 recruitment was not dependent on prior engagement with the health care system.[24-27,] Three of
- 24 the nine interventions took into account affordability of SACT, by reimbursing patients for taking part
- 25 in intervention, the rest six interventions did not mention direct or indirect costs associated with
- 26 participation in a study. [25,30,31]. Finally, all but four interventions attempted to incorporate the
- appropriateness of SACT services by increasing levels of engagement and promoting informed
 decision-making. [26,28,30-32] No intervention in this scoping review incorporated all five
- 29 dimensions of the equity framework. Language barriers were addressed in studies by Im et.al and Loi
- 30 et.al., as participants' native languages were Asian and Spanish, respectively.
- Table 3. Overview of included studies detailing interventions to improve experiences and clinical outcomes in
 people from MEG receiving SACT treatments
- 33
- 34
- 35
- 36 37
- 38
- 39
- 40

Author, year, country	Intervention delivery mode	Objectives	Methods	Sample Demographics	Outcomes	Quality assessment
Rust, 2015, USA ^[30]	In person Medication adherence skills training (MST) workshop	To explore what was the level and role of health literacy, with respect to medication adherence and self-efficacy.	Randomised pilot study	African American breast cancer survivors within one year of treatment from three urban areas of a Southeastern, Tennessee State (n=48)	Questionnaires were administered to measure patient self- efficacy in medication usage, medication adherence, and a three question measure for health literacy. Intervention did not show a statistically significant effect on health literacy (HL), medication adherence or self-efficacy. Statistically significant relationship was found between the initial HL and medication adherence.	3
Rosenzweig, 2011, USA ^[32]	Hybrid (in person and digital) Psychoeducational one to one intervention	To test the effect of an intervention on treatment adherence.	Randomised pilot study	African American breast cancer women receiving first adjuvant therapy from two urban sites of the Comprehensive Breast Program in the University of Pittsburgh Cancer Institute, a National Cancer Institute (n=24)	45-minute face-to-face meeting with intervention participants was recorded and adherence rates to chemotherapy were measured at three time points corresponding to treatment completion: 50%, 75%, and 100%. Results demonstrated more rapid initiation of chemotherapy and better overall adherence to chemotherapy.	2
Cykert, 2019, USA ^[26]	Digital Real time registry combined with feedback	To improve the treatment completion rates of surgery, recommended radiation and chemotherapy for each patient.	Non- randomised controlled trial (NRCT)	African American and White ethnic group patients with early stage lung and breast cancers at two cancer centres (n=302)	African American patients in the intervention group achieved a Treatment Complete rate of 88.4% compared to 89.5% for White ethnic group (p = 0.77).	1
Loi, 2016, USA ^[24]	Digital Self-administered stress management training	To examine the efficacy of a culturally and linguistically tailored training in improving QOL and reducing psychological distress.	Randomised controlled trial (RCT)	Hispanic/Latino patients newly diagnosed with cancer from 17 local community oncology practices across the U.S. and Puerto Rico (n=219)	No significant treatment effects on quality of life and reducing psychological distress were demonstrated. Improved mental health scores were observed with patients on a psychotropic agent (p=.04).	2

Turbes, 2015, USA ^[27]	Digital Web-Based Programme using interviews, online screener and post-use survey	To assess implementation and fidelity of an intervention.	Mixed- method study	African American women age <45 with breast cancer from three cancer centres (n=1442)	75% of post-use survey respondents were very or somewhat satisfied with the web-based programme; 70% of respondents said the web-based programme content was somewhat or very useful.	1
Sheppard, 2013, USA ^[31]	In person Peer-Led decision support intervention	To assess the acceptability of the intervention.	Non- randomised pilot study	African American women over 21 years old diagnosed with any stage breast cancer from Washington, DC metropolitan area (n=76)	Participants reported increased self-efficacy in communicating with providers (70 %) and self-efficacy in making treatment decisions (70 %).	2
Perez, 2020, USA ^[29]	Digital Interactive cancer- communication video program	To examine the feasibility and acceptability of an interactive video program.	Randomised pilot study	African American women newly diagnosed with breast cancer from three cancer centres (n=107)	104 of 108 patients allocated to the intervention reported moderate-to-high levels of positive emotional reactions to stories and identification with storytellers.	1
Thompson, 2021, USA ^[25]	Digital Viewing survivor stories	To determine whether viewing survivor stories improved newly diagnosed African American breast cancer patients' QOL.	Randomised controlled trial (RCT)	African American women with breast cancer from three cancer centres (n=228)	No effect of study arm on QOL, depressive symptoms, or concerns about recurrence was found in this study.	1
lm, 2023, USA ^[28]	Digital Virtual program using social media sites, interactive online educational sessions and online recourses	To determine the efficacy of a culturally tailored virtual information and coaching/support program in improving patients survivorship experience.	Randomised controlled trial (RCT)	Asian American women within 5 years of the diagnosis from online and offline cancer support groups (n=199)	Women were asked to fill out the questionnaires at different points during 12week study period. Results showed that intervention group had a significant increase in their quality of life.	2

1 *Key to quality assessments: 1-Good, 2-Average, 3-Poor

2 <u>Quality assessment/Critical appraisal of individual sources of evidence</u>

3 Only one study [30] did not meet MMAT criteria and therefore was scored as a 3. Another four

4 studies have met the criteria at 60-80%, thus were given a score of 2. The remaining four studies

5 have met the MMAT criteria at 80-100% and were rated as good quality studies with a score of 1.

2 Table 4. Accessibility dimensions identified in interventions targeting people in MEG receiving SACT^[5]

Author, year, country	Positive impacts (Intended and/or Unintended)	Negative impacts (Intended and/or Unintended)	Dimensions of accessibility
Rust, 2015, USA ^[30]	Approachability: Participants were recruited by the staff of a community- based organisation for underserved and minority women diagnosed with breast cancer, helping participants to engage and connect with the study personnel. Affordability: All participants were compensated for participation with gift cards at the beginning and end of the study, however costs of transportation or treatment were not discussed in the study. Appropriateness: Patients were able to engage with a licensed pharmacist and social worker during two-hour workshop exploring medication usage and adherence.	Acceptability: Translational services were not mentioned in the study. Only the possibility of an oral questionnaire delivery was suggested. Availability: Even though participants for this intervention were recruited from three urban locations, increasing it's availability to reach several communities, the sample size was small, included only 48 patients with a very specific set of characteristics, therefore reducing the chances of taking part in the intervention for those individuals who were beyond first year of breast cancer treatment path. Participants were recruited through contacts with American Cancer Society or through community-based organisation, limiting its reach to patients from other organisations.	Approachability ** Acceptability ** Availability ** Affordability ** Appropriateness **
Rosenzweig, 2011, USA ^[32]	Approachability: Race-matched recruiters were used during recruitment phase of the study. Acceptability: The interventionist was an African American breast cancer survivor. Appropriateness: Patients were provided with 1:1 supportive session discussing attitudes (including perceptions and stressors) that may affect adherence to clinical visits and treatment.	Availability: Patients were recruited from two University of Pittsburgh Cancer Institute clinics, thus limiting the participation in this intervention for wider communities. Affordability: Consideration of cost was mentioned only as limitation in this study. Cost to retain an interventionist on staff needs to be assessed to determine whether the intervention can be integrated into routine clinical practice.	Approachability ** Acceptability ** Availability ** Affordability ** Appropriateness **
Cykert, 2019, USA ^[26]	Approachability: multi-faceted approach utilizing transparency of clinical data and care team accountability achieved through race-specific audit and feedback was required for the highest probability of success. Acceptability: Nurse and physician were specially trained in teach - back technique, anti-racism, and to advocate for patients. Availability: Study spanned over 5 years and nurse navigators have applied their special anti-racial training to all patients. Appropriateness: all patients were engaged in the study by trained personnel	Affordability: Cost implications of treatment or transportation were not mentioned, apart from a grant from The National Cancer Institute that was used to fund the study.	Approachability ** Acceptability ** Availability * Affordability ** Appropriateness **
Loi, 2016, USA ^[24]	Approachability: patients received targeted intervention Acceptability: Community experts and a certified translator were consulted for the initial adaptation and translation of English materials. Availability: patients were recruited from 17 community practices, thus expanding the pool of participants significantly.	Affordability: The costs of treatment or transportation were not accounted in the study. Appropriateness: 5-minute, standardized explanation of the nature and purpose of the intervention was provided. Patients were instructed to view the video/DVD first and to follow the directions in the booklet to better understand the training, practice and use of stress management techniques.	Approachability ** Acceptability ** Availability * Affordability * Appropriateness *

Turbes, 2015, USA ^[27]	Approachability: Participants received reproductive and psychosocial information and support using various platforms during the study. Acceptability: Materials of the intervention were culturally appropriate for the target population. Availability: intervention was provided via multiple channels, including web and social	Affordability: Ability to pay indirect costs such as transportation not covered Appropriateness: This intervention was highly reliant on technology, assuming that participants have access to internet and social media. Also this type of intervention does not engage patients to high extent in the decision making about their treatment.	Approachability ** Acceptability ** Availability ** Affordability ** Appropriateness **
Sheppard, 2013, USA ^[31]	media. Approachability: Women with histologically confirmed breast cancer were recruited from the Washington, DC area Acceptability: The coach used a culturally appropriate guidebook and decision- making model—TALK Back! Appropriateness: Peer-delivered culturally relevant decision support intervention for Black women with breast cancer. Affordability: Ability to pay indirect costs such as transportation not covered. Participants received a grocery store gift card.	Availability: Patients were recruited mainly from cancer surgeons.	Approachability ** Acceptability ** Availability ** Affordability ** Appropriateness **
Perez, 2020, USA ^[29]	Approachability: Intervention arm completed a baseline/pre-intervention interview, received the video intervention, and completed a post-intervention 1- month follow-up interview. Acceptability: Intervention was conducted by specially trained study team coordinators.	Availability: Patients were recruited from their breast surgeons Affordability: Ability to pay indirect costs such as transportation not covered Appropriateness: Patients received a brief (~10 minute) in-person training to use the video program plus an instructional user guide to take home.	Approachability ** Acceptability ** Availability ** Affordability ** Appropriateness **
Thompson, 2021, USA ^[25]	Approachability: African American women with non-metastatic breast cancer interviewed five times over two years. Acceptability: The video used in the intervention was culturally adapted to the participants. Availability: Participants of the study were interviewed five times over a period of two years, intervention was reliant on ability to use tablet or computer Affordability: Ability to pay indirect costs such as transportation not covered. Participants received \$25 per interview.	Appropriateness: Patients were instructed to watch videos and understand and interpret their content in their own homes.	Approachability ** Acceptability ** Availability ** Affordability ** Appropriateness **
Im, 2023, USA ^[28]	Approachability: Support programme was created for Asian-American women with breast cancer Acceptability: The content of interventional material was culturally tailored to Asian- American women (intervention components that were provided in five languages (English, Mandarin Chinese [Simplified and Traditional], Korean, and Japanese) Appropriateness: The social media sites provided a medium by which participants could connect to each other and share their own breast cancer survivor experience with peers.	Availability: The intervention group utilized American Cancer Society [ACS] website to deliver intervention Affordability: Ability to pay indirect costs such as transportation not covered	Approachability ** Acceptability ** Availability ** Affordability ** Appropriateness **

1 <u>Discussion</u>

2 This was the first scoping review of its kind investigating interventions to optimise SACT treatments in 3 people with cancer from MEG. The review showed the breadth of interventions that have been 4 researched that could have the potential to improve patient experience, self-efficacy when making 5 decisions related to SACT and adherence. Our searches found only a small number of studies in total, 6 with differing outcome measures. Studies developed were focussed on digital interventions that 7 were either offering holistic support to patients or educational materials. Comparing digital versus in-8 person delivered interventions showed that there was no difference in one mode of delivery 9 superseding another, where equity related provisions were considered. Interventions that were most 10 effective in improving access to SACT used variety of approaches, ensuring the design of the study 11 integrated wide range of accessibility parameters, such as approachability, acceptability, availability, 12 affordability, and appropriateness of the studies. We adapted an existing equity framework [6] to 13 assess patients' from MEG experiences during their SACT treatments. Our results suggested that 14 structural and social determinants of health were interconnected and highlighted the importance of 15 incorporating these equity considerations when developing interventions. It was surprising that 16 none of the studies were conducted outside of the USA. The UK population is 18% from MEG and 17 therefore inclusive cancer care would benefit 1 in 5 patients. [33] Another unexpected finding was 18 that none of the studies had assessed the economic impact of interventions, which is essential for 19 health care researchers, policy makers and providers to make informed decisions. 20 The majority of studies were describing interventions in early breast cancer – a finding that is not 21 surprising as breast cancer is among the top three cancers globally, affecting 2.26 million women and 22 is the leading cause of cancer mortality worldwide [34]. There are multiple factors behind disparities 23 that are affecting global cancer survival rates and it is one of the World Health Organisations' (WHO) 24 priorities to improve survival across the world. Breast cancer patients are being offered a 25 comprehensive treatment and supportive care, but still adherence issues are very common and

- 26 understood to have health economic implications [35]. Non-adherence to prescribed medications is
- associated with poor therapeutic outcomes, progression of disease and overall healthcare costs.
- 28 Improvement in adherence can help in reducing the economic burden in long-term and improve
- cost-effectiveness [36]. Significant economic barriers to cancer services still exist and people from
 MEG are impacted by structural vulnerability, combining factors such as poverty, homelessness and
- 31 racism. Research indicates the populations that experience socioeconomic disadvantage more likely
- 32 to experience delays in starting SACT treatment, are less likely to receive any treatment and have
- 33 poor adherence rates to systemic therapy. Interestingly, results of this scoping review show that
- 34 digital interventions were chosen as a preferred method of conveying information to people from
- 35 MEG, suggesting that there is access to technology and capacity to be able to self-manage their care
- 36 needs and navigate a complex care system. [37]
- 37 Following the Covid-19 pandemic there have been an increased number of virtually delivered
- 38 healthcare interactions [38,39] and they are still being used daily in current practice worldwide,
- 39 albeit mostly in high-income countries. Digital interventions reduce costs associated with
- 40 appointment scheduling and can reduce travel time for patients to cancer centres, thus translating
- 41 into increased patient satisfaction with their care. Virtual coaching and digital support programmes
- 42 could potentially change health behaviours and as a result improve health outcomes amongst
- 43 vulnerable cancer patients. However, using telehealth as a means of providing healthcare services to
- 44 people with cancer from MEG could be also seen as another potential avenue for widening inequity
- 45 among cancer patients, as low and middle-income countries should be able to benefit from it too
- 46 [40].

- 1 One previous systematic review by Hayanga et. al [41] suggested that for some people from MEG
- 2 with multiple long-term conditions in the United Kingdom there may be inadequate initiatives for
- 3 managing health conditions and that there is a need for enhanced strategies to reduce ethnic
- 4 inequalities in healthcare. Wider societal processes such as suboptimal healthcare provision together
- 5 with individual cancer patients' from MEG journey (with additional barriers, such as language
- 6 difficulties and poor literacy skills) can have negative impact on people's ability to access and utilise
- 7 healthcare services effectively.
- 8 Another study by Unruh et.al, comparing health policy responses to COVID-19 highlighted that there
- 9 needs to be a universal investment across the health sector physical infrastructure and training to
- 10 reduce unmet care needs and health inequalities among the most vulnerable population groups.
- 11 [42]. Interventions to reduce health inequity should be directed at downstream determinants of
- 12 health, such as individual health-care needs, midstream determinants, such as neighbourhood
- 13 conditions, or upstream determinants, such as structural racism and discrimination. There are several
- 14 theoretical approaches to social determinants of health with ethnicity and racism falling under a
- 15 social disadvantage approach. Greater social disadvantage is associated with poorer health and more
- 16 research is needed to clarify the underlying pathways.
- 17 This study has identified gaps in the design of interventions targeted at people from MEG
- 18 populations and a new equity-framework was tested for patients undergoing SACT. Future
- 19 interventions exploring the relationship among minority patient groups may use this framework as a
- 20 useful tool in their equity assessments to improve access to SACT across all at-risk groups and reduce
- 21 inequalities in cancer care.

22 <u>Study Limitations</u>

23 This is the only study of its kind that we are aware of, utilising a systematic methodology including 24 quality appraisal by two researchers, incorporation of an equity-based framework and intervention 25 reporting guidelines. Despite this, there were very few studies that met our inclusion criteria, and we 26 were limited by small sample sizes. By nature, scoping reviews capture the breadth of literature in 27 the area and therefore our studies were heterogeneous in the methods used. This scoping review 28 maps out the currently available literature on interventions in people with cancer from MEG and 29 highlights the need for further adequately powered interventional studies. Future work should focus 30 on further evaluation of SACT services provided in cancer centres, starting with a local context 31 assessment and review of policies.

32 <u>Conclusions</u>

33 The key findings were that published evaluations of interventions to optimise SACT management in 34 people from MEG are limited to early breast cancer patients, predominantly of African-American 35 background, and that studies were mainly conducted in high-income countries, such as USA in 36 particular. We found evidence of interventions incorporating the equity domains that reported 37 improved patient engagement and experience. This new knowledge will help to implement future 38 SACT interventions, addressing health inequities across the cancer continuum and explore future 39 directions in patient navigation to promote equitable care. Healthcare providers should view the 40 equity considerations identified in this review as unique perceptions of people from MEG undergoing 41 SACT treatment rather than obstacles in providing cancer care. The future research should focus on 42 psychosocial and cultural influences to help design effective equitable interventions.

1		
2		References
3 4 5	1.	Torre LA, Siegel RL, Ward EM, Jemal A. Global cancer incidence and mortality rates and trends—an update. Cancer epidemiology, biomarkers & prevention. 2016 Jan 1;25(1):16-27. DOI: 10.1158/1055-9965.EPI-15-0578
6	2.	Singh GK, Jemal A. Socioeconomic and racial/ethnic disparities in cancer mortality, incidence,
7		and survival in the United States, 1950–2014: over six decades of changing patterns and
8		widening inequalities. Journal of environmental and public health. 2017
9 10	2	Oct;2017. DOI: 10.1155/2017/2819372 Patel ML Forguson IM, Castro F, Poreira Ectromora CD, Armaiz Boña GN, Duron V, Hlubocky F,
10	5.	Infantado A. Nuqui B. Julian D. Nortev N. Racial and ethnic disparities in cancer care during
12		the COVID-19 pandemic. JAMA network open. 2022 Jul 1;5(7):e2222009
13		DOI: <u>10.1001/jamanetworkopen.2022.22009</u>
14	4.	Patel MI, Lopez AM, Blackstock W, Reeder-Hayes K, Moushey EA, Phillips J, Tap W. Cancer
15		disparities and health equity: a policy statement from the American Society of Clinical
16		Oncology. Journal of Clinical Oncology. 2020 Oct 10;38(29):3439-48.
17	-	DOI: <u>10.1200/JCO.20.00642</u>
18 10	5.	sayani A, Ali MA, Dey P, Corrado AM, Ziegler C, Nicholson E, Lotters A. Interventions designed
20	6	Szczenura A Price C Gumber A Breast and howel cancer screening untake patterns over 15
21	0.	vears for UK south Asian ethnic minority populations, corrected for differences in socio-
22		demographic characteristics. BMC public health. 2008 Dec;8:1-5. doi: 10.1186/1471-2458-8-
23		346.
24	7.	Clinical and Research Reports. 2023 Mar 1;4(3):100469. DOI: <u>10.1016/j.jtocrr.2023.100469</u>
25		Trends in the global medicines market 2024. GHP PDIG Symposium, 20th June 2024. IQVIA
26		Institute for Human Data Science. Global Outlook to 2028, May 2024. Available from:
27	•	www.iqviainstitute.org. Accessed [17/9/2024]
28	8.	Greer JA, Amoyal N, Nisotel L, Fishbein JN, MacDonald J, Stagi J, Lennes I, Temei JS, Safren
29 30		oncologist 2016 Mar 1:21(3):354-76
31	9	McGuinness S. Hughes L. Moss-Morris R. Hunter M. Norton S. Moon 7. Adherence to
32	0.	adjuvant endocrine therapy among White British and ethnic minority breast cancer
33		survivors in the United Kingdom. European Journal of Cancer Care. 2022 Nov;31(6):e13722.
34	10	. Cancer incidence statistics. Available at: < Cancer incidence statistics Cancer Research UK> Accessed on
35		22.04.24
36	11.	. Verbrugghe M, Verhaeghe S, Lauwaert K, Beeckman D, Van Hecke A. Determinants and
3/ 20		associated factors influencing medication adherence and persistence to oral anticancer
30 39		DOI: 10.1016/j.ctry 2012.12.014
40	12	Papadakos IK, Hasan SM, Barnsley I, Berta W, Fazelzad R, Papadakos CI, Giuliani MF, Howell
41		D. Health literacy and cancer self-management behaviors: A scoping review. Cancer. 2018
42		Nov 1;124(21):4202-10. DOI: <u>10.1002/cncr.31733</u>
43	13	. Gast A, Mathes T. Medication adherence influencing factors—an (updated) overview of
44		systematic reviews. Systematic reviews. 2019 Dec;8:1-7. DOI: <u>10.1186/s13643-019-1014-8</u>
45	14	. Berkman ND, Davis TC, McCormack L. Health literacy: what is it?. Journal of health
46		communication. 2010 Aug 31;15(S2):9-19. DOI: <u>10.1080/10810730.2010.499985</u>
47	15	. Hyde JE. Reconsidering Health Literacy: The Role of Implicit Bias. Temple University; 2017.

1	16.	Vela MB, Erondu AI, Smith NA, Peek ME, Woodruff JN, Chin MH. Eliminating explicit and
2		implicit biases in health care: evidence and research needs. Annual review of public health.
3		2022 Apr 5;43(1):477-501. DOI: <u>10.1146/annurev-publhealth-052620-103528</u>
4	17.	Vaccarella S, Georges D, Bray F, Ginsburg O, Charvat H, Martikainen P, Brønnum-Hansen H,
5		Deboosere P, Bopp M, Leinsalu M, Artnik B. Socioeconomic inequalities in cancer mortality
6		between and within countries in Europe: a population-based study. The Lancet Regional
7		Health–Europe, 2023 Feb 1:25, DOI: 10.1016/i.lanepe,2022.100551
8	18	Arksey H. O'malley L. Scoping studies: towards a methodological framework. International
9		iournal of social research methodology 2005 Feb 1:8(1):19-32 doi
10		10 1080/1364557032000119616
11	10	Tricco AC Lillie F. Zarin W. O'Brien KK. Colguboun H. Levac D. Moher D. Peters MD. Horsley T.
12	10.	Weeks L. Hempel S. PRISMA extension for sconing reviews (PRISMA-ScR): checklist and
12		evolution Annals of internal medicine 2018 Oct 2:169(7):467-73 DOI: 10.7326/M18-
10		0850
14	20	NAL Pact Practice, How to clarify a clinical question [Internet] [cited 2024 Jan 15] Available
16	20.	from: https://bestpractice.bmi.com/info/us/toolkit/learn-ebm/how-to-clarify-a-clinical-question/
17	21.	McDonagh M, Peterson K, Raina P, Chang S, Shekelle P. Avoiding bias in selecting studies.
18		Methods guide for effectiveness and comparative effectiveness reviews [Internet]. 2013 Feb
19		20. PMID: 23487864
20	22.	Hoffmann TC. Glasziou PP. Boutron I. Milne R. Perera R. Moher D. Altman DG. Barbour V.
21		Macdonald H. Johnston M. Lamb SE. Betterof interventions: template for intervention
22		description and replication (TIDieR) checklist and guide. Bmi. 2014 Mar 7:348. doi:
23		10.1136/bmi.g1687 (www.equator-network.org/reporting-guidelines/tidier/)
24	23	Hong ON Eabregues S. Bartlett G. Boardman F. Cargo M. Dagenais P. Gagnon MP. Griffiths F.
25	20.	Nicolau B. O'Cathain A. Bousseau MC. The Mixed Methods Appraisal Tool (MMAT) version
26		2018 for information professionals and researchers. Education for information, 2018 Jan
20 27		1·34(4)·285-91 doi 10 3233/FEI-180221
28	24	Loi CX Nesman TM Xu P Taylor TR McMillan S Krischer IP Tyc VI. Gross-King M Huegel V
20 29	27.	A self-administered stress management intervention for hispanic nations undergoing
20		cancer chemotherany, Journal of Immigrant and Minority Health, 2017 Oct-19:1121-31
21		DOI: 10, 1007/c10002-016-0524
30	25	Thompson T. Bérez M. Van V. Kreuter MW/ Margenthaler IA. Colditz GA. Jeffe DB
3Z 22	20.	Pandamized controlled trial of a broast cancer Surviver Stories intervention for African
33 24		American warman Social Science & Medicine, 2021 Sch 1:270:112662
34 25		American women. Social Science & Medicine. 2021 Feb 1,270.115005.
35 26	26	DUI: 10.1016/J.SOCSCIME0.2020.113663
30 07	20.	Cykert S, Eng E, Manning MA, Robertson LB, Heron DE, Jones NS, Schaal JC, Lightroot A,
37		Zhou H, Yongue C, Giziice Z. A multi-faceted intervention aimed at Black-white disparities in
38		the treatment of early stage cancers: the ACCORE pragmatic quality improvement trial.
39		Journal of the National Medical Association. 2020 Oct 1;112(5):468-77.
40	07	DUI: <u>10.1016/j.jnma.2019.03.001</u>
41	27.	Johnson-Turbes A, Schlueter D, Moore AR, Buchanan ND, Fairley TL. Evaluation of a web-
42		based program for African American young breast cancer survivors. American journal of
43		preventive medicine. 2015 Dec 1;49(6):S543-9. DOI: <u>10.1016/j.amepre.2015.09.003</u>
44	28.	Im EO, Chee W, Paul S, Choi MY, Kim SY, Deatrick JA, Inouye J, Ma G, Meghani S, Nguyen GT,
45		Schapira MM. A randomized controlled trial testing a virtual program for Asian American
46		women breast cancer survivors. Nature Communications. 2023 Oct 14;14(1):6475.
47		DOI: <u>10.1038/s41467-023-42132-6</u>

1	29.	Pérez M, Kreuter MW, Yan Y, Thompson T, Sefko J, Golla B, Margenthaler JA, Colditz G, Jeffe
2		DB. Feasibility and acceptability of an interactive cancer-communication Video program
3		using African American breast cancer survivor stories. Journal of health communication.
4		2020 Jul 2:25(7):566-75. DOI: 10.1080/10810730.2020.1821132
5	30	Rust CE, Davis C. Moore MR. Medication adherence skills training for African-American
6	00.	breast cancer survivors: the effects on health literacy medication adherence, and self-
7		officacy. Social work in health care, 2015 Jan 2:54(1):22.46
/ 0		D_{1} 10 1020/00021220 2014 064447
0	24	DUI. <u>10.1080/00981589.2014.904447</u>
9	31.	Sneppard VB, waiiington SF, willey SC, Hampton RW, Lucas W, Jennings Y, Horton S, Muzeck
10		N, Cocilovo C, Isaacs C. A peer-led decision support intervention improves decision
11		outcomes in black women with breast cancer. Journal of Cancer Education. 2013
12		Jun;28:262-9. DOI: <u>10.1007/s13187-013-0459-z</u>
13	32.	Rosenzweig M. The attitudes, communication, treatment, and support intervention to
14		reduce breast cancer treatment disparity. Number 1/January 2011. 2011 Jan 1;38(1):85-9.
15		DOI: <u>10.1188/11.ONF.85-89</u>
16	33.	Ethnicity facts and figures. Goverment data about the UK's different ethnic groups. Available
17		at: < Ethnicity facts and figures – GOV.UK (ethnicity-facts-figures.service.gov.uk)> [Accessed
18		on 24/09/2024].
19	34.	Ferlay J, Colombet M, Soerjomataram I, Mathers C, Parkin DM, Piñeros M, Znaor A, Bray F.
20		Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and
21		methods. International journal of cancer. 2019 Apr 15;144(8):1941-53.
22		DOI: <u>10.1002/ijc.31937</u>
23	35.	Wilkinson L, Gathani T. Understanding breast cancer as a global health concern. The British
24		journal of radiology. 2022 Feb 1;95(1130):20211033. DOI: <u>10.1259/bjr.20211033</u>
25	36.	Gupta J, Joshi P, Kamra S, Sehgal M. Economic Burden due to Treatment Non-Adherence in
26		Patients with Breast Cancer: A Systematic Review. Value in Health. 2016 May 1;19(3):A156.
27		Doi: 10.106/jval.2016.03.1461
28	37.	Bourgeois A, Horrill TC, Mollison A, Lambert LK, Stajduhar KI. Barriers to cancer treatment
29		and care for people experiencing structural vulnerability: a secondary analysis of
30		ethnographic data. International Journal for Equity in Health. 2023 Mar 30;22(1):58.
31		DOI: <u>10.1186/s12939-023-01860-3</u>
32	38.	Chow MC, Chambers P, Singleton G, Patel J, Cooper S, Mythen C, Bautista-González E,
33		Chisnall G, Djellouli N, Thwaites B, Wong IC. Global changes to the chemotherapy service
34		during the covid-19 pandemic. Journal of Oncology Pharmacy Practice. 2021 Jul;27(5):1073-
35		9. DOI: <u>10.1177/10781552211015767</u>
36	39.	Dalby M, Hill A, Nabhani-Gebara S. Cancer patient experience of telephone clinics
37		implemented in light of COVID-19. Journal of Oncology Pharmacy Practice. 2021
38		Apr;27(3):644-9. DOI: <u>10.1177/1078155221990101</u>
39	40.	Doraiswamy S, Abraham A, Mamtani R, Cheema S. Use of telehealth during the COVID-19
40		pandemic: scoping review. Journal of medical Internet research. 2020 Dec 1;22(12):e24087.
41		DOI: <u>10.2196/24087</u>
42	41.	Hayanga B, Stafford M, Becares L. Ethnic inequalities in healthcare use and care quality
43		among people with multiple long-term health conditions living in the United Kingdom: a
44		systematic review and narrative synthesis. International Journal of Environmental Research
45		and Public Health. 2021 Nov 29;18(23):12599. doi: 10.3390/ijerph182312599.
46	42.	Unruh L, Allin S, Marchildon G, Burke S, Barry S, Siersbaek R, Thomas S, Rajan S, Koval A,
47		Alexander M, Merkur S. A comparison of 2020 health policy responses to the COVID-19
48		pandemic in Canada, Ireland, the United Kingdom and the United States of America. Health
49		Policy. 2022 May 1;126(5):427-37. DOI: <u>10.1016/J.healthpol.2021.06.012</u>

2 Figure 1. Equity-orientated SACT intervention considerations for patients from MEG.

Approachability	Acceptability	Availability	Affordability	Appropriateness
Ability to perceive	Ability to seek	Ability to reach	• Ability to pay	Ability to engage
Access to	SACT related	Transportation/	Cost of	Patients are not
hotline	information/mat	Mobility	treatment	left to receive, understand and
Knowledge	erials is tailored	Ability to	Eligibility for free	act on
about support	cultural values	access/use	treatment	information
services		technology	Carer	provided
Personalized	Translator	Community	responsibility	Trained
risk assessment	provided	engagement		coordinators
		Intervention	Childcare	Informed
Targeted health	Sociocultural	provided via	Reimbursement	decision-
campaign	training for staff	multiple	for participation	making
ourrpaiBit		channels		encouraged

3

1

4 Figure 2. Prisma flow diagram

